

# CHARACTERISTICS OF NUTRIENTS IN THE ESTUARIES OF KANYAKUMARI DISTRICT-TAMILNADU-INDIA.

Dr.kokila.p1 and Rathika.s2

1Head of the department,Vivekanandha college of arts and science, Kanyakumari  
2Department of science and Humanities,Rohini college of engineering and technology  
Kanyakumari District-TAMILNAD, krhanshika@gmail.com


**ABSTRACT:** Water samples were collected in four major estuaries of Kanyakumari District to analyse the distribution of dissolved nutrients and it is said to be polluted. Water samples were analysed for physico-chemical parameters including pH, turbidity, electrical conductivity, total hardness, dissolved oxygen, BOD and anions were analysed like Ca, Mg, Fe, free ammonia, NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, PO<sub>4</sub><sup>2-</sup>, F<sup>-</sup> and Cl<sup>-</sup> during the different seasons of monsoon period from August - January (2011-2012). The concentrations of most of the investigated parameters in the water sample from Manakudy (Station 1), Rajakamangalam (Station 2), Kadiyapattanam (Station 3), Thengapattanam (Station 4) estuaries in Kanyakumari district were exceeded the permissible limit of WHO, CPHEEO water quality guidelines.

**Keywords:** Estuary, Nitrate, Dissolved Oxygen, WHO, CPHEEO.



## Introduction

Rivers are the main inland water resources for domestic, industrial and irrigation purposes and often carry large municipal sewage, industrial waste water discharges and seasonal runoff from agricultural land to the coastal region. It is for this reason that the river water is mostly enriched in nutrients compared to other environments. The four major rivers that flow in Kanyakumari district are, Thambaraparani, Valliyar, Pazhayar and Paraliyar. Thambaraparani river confluences with Arabian sea near Thangaipattanam estuary about 56 km, west of Kanyakumari District. The Valliyar river collects the drainage from P.P. Channel and its branches ayacuts and confluences with Arabian sea in Kadiyapattanam estuary. The Pazhayar river collects the drainage of Thovalai, Ananthanar and N.P. and confluences with Arabian sea in Manakudy estuary. The Paraliyar flows through the mathur hanging trough which is the highest and longest aqueduct in Asia and confluences with Arabian sea in Rajakkamangalam estuary. Usually in the near shore water and estuaries exhibit considerable seasonal variations depending on the local conditions of rainfall, tidal incursions, quantum of fresh H<sub>2</sub>O inflow, affecting the nutrient cycle of different coastal environments (Choudury & Panigraphy 1991). Hence the present study is designed to investigate the changes of physico-chemical parameters and the nutrients in the above mentioned estuaries.



**Fig. 1-Map of Estauries in Kanyakumari district.showing study sites(st.1- Manakudy(D),st.2- Rajakamangalam(c),st.3-Kadiyapattanam(B),(st.4-Thengapattanam(A )**

## Materials & Methods

Water samples were collected in the different seasons of monsoon period from Aug. 2011-Jan.2012. Surface water samples were collected in a clean plastic bucket. Preservation and transportation of the water samples to the laboratory were as per the standard methods (APHA 1998).The samples were analysed in 15 different parameters. PH was measured using a PH analyzer (WTW model multi 340 ) while conductivity was analysed by Mohr-Kundensen  $\text{AgNO}_3$  titration method (APHA,1998) .Turbidity was measured by Nephelometer using 0.02 NTU standards. Dissolved oxygen was fixed immediately after collection and then determined by winkler's method. Nutrients ( $\text{NH}_4^+$ , $\text{NO}_3^-$ ,  $\text{PO}_4^{3-}$ ,Fe) were determined by standard photometric method. (Grasshoff, et 1999) using varian 50 bio U.V visible spectrophotometer. Ions  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and hardness were determined by the complexometric EDTA titration method using EBT as a indicator (vogel,1978). samples for BOD were incubated in laboratory for five days at  $20^\circ\text{c}$  (Trivedy and Goel, 1984) . Whatman 541 filter papers were used for the determination of total dissolved solids.The present study reports the seasonal pattern of the physico-chemical parameters at these four important estauries.

## Results & Discussion

In this study analysis, water quality in different seasons were carried out to determine the seasonal variation of physical and chemical characteristic of water. The results were listed in table 1, and fig. - 1,2. The appearance of water sample and its odour are noted. Water colour during the north east monsoon period is brownish and algae odour due to the rainfall and sediments flow.

PH of the estaurine water was alkaline and fluctuated between 7.16 - 8.57. It was well within the limits (ICMR standard; 7-8.5). Maximum values were noted at station 3 during the north east monsoon period and minimum was recorded at station 2 during the post monsoon period. Electrical conductivity of estaurine was varied in the range 233 - 16986 mics/cm. The average conductivity was 5346 mic s/cm. The conductivity value was very high at station 1, during the monsoon & post monsoon period due to high salinity. This is because of wide variation in the discharge of river water.(Mudali Bipra et al.) Turbidity is mainly due to the dispersion of suspended particles. Turbidity values ranged from 2-41 NTU. Maximum was recorded at station 3, during the post monsoon period and minimum was recorded at station 4 during Monsoon and north east monsoon periods.

**Table1:Physico chemical analysis of water in different seasons**

seasons	stations	Physical parameters			
		Turbidity Units	NT	Total Solids[mg/lit]	Dissolved Electrical Conductivity[micS/cm]
Monsoon [August,September]	1	4		10482	15881
	2	2		3273	4959
	3	1		247	374
	4	2		2918	4421
North East Monsoon October,November]	1	13		722	1078
	2	17		2227	3324
	3	13		263	393
	4	2		6223	9430
Post Monsoon [December,January]	1	34		11013	16986
	2	32		1124	1703
	3	41		154	233
	4	35		3546	5373

seasons	statio ns	Chemical parameters											Po <sub>4</sub> <sup>2-</sup>	DO	BOD
		PH	Hard ness	Ca <sup>2+</sup>		Fe	NH <sub>4</sub>	NO <sub>3</sub>	Cl <sup>-</sup>	F <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>				
Monsoon													-		

[August,September]					Mg <sup>2+</sup>									
	1	7.57	610	640	1080	0.24	0.86	2	7200	0.4	74	1.6	0	15
	2	7.22	600	60	108	0.24	2.18	4	800	0.4	106	0.7	6.4	21
	3	8.16	30	8	2	0.47	0.46	2	20	0.4	4	0.40	7.3	12
	4	6.32	800	40	168	0.24	1.644	4	1350	0.2	171	0.40	6.7	18
North East Monsoon	1	7.54	2020	152	394	0.24	1.12	7	2800	0.4	429	0.80	1.4	32
	2	7.88	0	3120	136	0.24	1.46	9	2100	0.4	343	0.75	1.2	36
October,November]	3	8.57	40	8	5	0.12	1.08	4	44	0	7	0.90	1.3	28
	4	7.32	2120	112	442	0.12	0.92	7	3100	0.2	286	0.7	1.3	28
42Post Monsoon	1	7.48	4600	720	672	3.06	0.88	10	6050	0.8	54	1.9	5.9	42
	2	7.16	640	60	118	0.12	0.23	9	1360	0.4	46	2.75	5.3	36
[December, January]	3	8.55	88	24	7	0.35	0.23	4	40	0.2	19	0.80	6.1	42
	4	7.68	790	68	149	0	0.08	9	1505	0.2	46	2.3	6.2	32

Dissolved Oxygen (DO) levels varied from 1.2 - 7.3 mg/l. Maximum value was observed at station 3, in monsoon period and minimum was observed at station 2 in north east monsoon period. This low value is due to the addition of high organic contents in the fishing jetty leading to oxygen depletion. BOD was observed to be in the range from 12 - 42 mg/l. Maximum value was observed at station 2 during post monsoon period and low value was recorded at station3 during monsoon period. The maximum values are due to the maximum biological affinity at elevated temp (Ghauzan, et 2006) and reduced flow of riverine water. Due to the presence of high calcium and magnesium content observed at station 1, during the north east monsoon period, the estuarine water exhibit high hardness. The minimum value of hardness was observed at station 3 during monsoon period. The nitrate (NO<sub>3</sub><sup>-</sup>) concentration ranged from 2 -10 mg/l. Maximum concentration was observed at station2 during post monsoon and minimum was observed at station 3 during monsoon period. This is due to domestic sewage, agricultural wash off, decaying plant, animal materials, fertilizers and other waste effluents containing nitrogenous compounds (Adeyeye and Abulude, 2004). Such source of nitrate was also reported from Mahanadi estuary (Sunderay, et. 2006).

Ammonium concentration was observed in the range from 0.08 - 2.18 mg / l. The maximum was observed at station 2 during the monsoon period. The minimum was observed at station 4 during post monsoon period. This low value is due to the influence of industrial wastes. Chloride concentration was observed the range from 20-7200 mg/l. The maximum was observed at station 1 during the monsoon period. The minimum was noted at station 3 during the monsoon period. The WHO standard desirable limit of chloride was 200 mg /l and permissible limit 600 mg/l. The observed values are exceeded the desirable

and permissible limit of WHO. High chloride content impacts taste and could cause corrosion (WHO). Iron concentration was observed the range from 0– 3.06. The maximum value was observed in station1 during the post monsoon period. Minimum value was noted at station 4 during the post monsoon period. CPHEEO standard desirable limit of iron was 0.1 and the permissible limit was 1. But the maximum observed value exceeded the permissible limit






Fig -1 .Distribution of physical nutrients at stations 1 to 4 of kanyakumari estauries during August - January (2011-2012)


Station-1



Station -2



Station-3



Station4




Fig -2 Distribution of nutrients at stations 1 to 4 of kanyakumari estuaries during August - January (2011-2012)

Fluoride and Sulphate levels were in the range of 0 - 0.8 mg/l and 4-429 mg/l respectively. Based on the CPHEEO standard fluoride desirable level should be 1 and the standard permissible limit 1.5 mg/l. WHO standard showed highest desirable limit of sulphate 200 mg /l and the maximum permissible limit should 400 mg/l.

## ACKNOWLEDGEMENT

The authors acknowledge with gratitude the help and support from Ms.Ramanibai and my friends for encourage to finish the project.

## CONCLUSION

Due to different environmental conditions and ecosystem, water quality standards vary significantly. The presence of trace metal contents in all the soil and crops, geographical location, pesticides, fungicides, fertilizers in the runoff water reached at the estuarine region has been polluted. The physico-chemical characteristics of estuarine water in the study area suggested that there was no harmful chemical contamination. But due to the exceeded value of chloride, iron, sulphate content may cause the corrosion which leads to the crackages in the bridges in the estuarine area. If proper measures are taken for the treatment of sewage before discharge and restrictions are put on various anthropogenic activities upstream, the estuary would remain healthy in the long run.

## REFERENCES

- 1) Adeyeya EI and Abulude Fo, *Analytical assessments of some surface and ground H<sub>2</sub>O resources in Ile - Ife, Nigeria*. J.Chem Soc. Nig. **29**: PP (98 - 103). **(2004)**.
- 2) APHA **(1998)**. Standard methods for the Examination of Water and Waste Water, American Public Health Association, Washington DC, USA.
- 3) Choudhury S.B & R.C Panigraphy, *Seasonal distribution and behaviour of nutrients in the greek and coastal water of Gopalpur, east Coast of India*: Maha Sagar Bull, Nati. Inst. Oeanogr ; **24(2)** : (91 - 88).**(1991)**.
- 4) Ghavzan N.J., Gunala, V.R., and Trivedy, R.K., (2006). *Limnological evaluation of an urban fresh water river with special reference to phytoplankton*. Pollution research; **25** PP ( 259-268). **(2006)**.
- 5) Grassshoff, K., Ehrhardt, M. and Kremling, K.(1999) .Methods of seawater analysis. pp(159-226) **.(1999)**
- 6) Muduli Bipra Prasanna, Panda Chitta Ranjan; *Physico chemical properties of water collected from Dharma estuary*. Environmental Science Vol **No.3**. **(2010)**.
- 7) Sunderay, S.K., Panda U.C., Nayak, B.B., and Bhatta, D., . Multi variate statistical techniques for the evaluation of spatial and temporal variation in water quality of Mahanadi river- estuarie system (India) a case study. Environmental Geochemistry and Health **28**: PP (317-330). **(2006)**.
- 8) Trivedy ,R.K.,and Goel,P.K.).Chemical and Biological methods for water pollution studies.ENV.PUB.Karad,India: pp(.104),**(1984)**
- 9) Vogel AI, A text book of Quantitative Inorganic analysis including Elementary Instrumental Analysis 4<sup>th</sup> Ed. The English Language book society and langman.
- 10) WHO, Guidelines for water quality, **vol.2** Health Criteria and other supporting information, Genera; WHO. **(1990)**