

**PHYTOCHEMICAL SCREENING AND ANTIMICROBIAL ACTIVITY OF THE  
ROOTS AND STEM BARK of Mahogany (*Khaya senegalensis*) and *Anogeissus leiocarpus*  
Grown in Bauchi Metropolis of Bauchi State, Nigeria.**

**Authors: <sup>1</sup>Yusuf H., <sup>2</sup>Bello K.S., <sup>3</sup>Ebu B., <sup>4</sup>Ijebor F.D.**

**<sup>1,2,3&4</sup> Federal Polytechnic, Bauchi. Department of Science Laboratory Technology.**

**Corresponding Author: Yusuf H [haleesadee5@gmail.com](mailto:haleesadee5@gmail.com) +2347034574714**

**ABSTRACT**

The roots and stem barks of *Khaya senegalensis* was prepared by extracting 100<sub>g</sub> of the crude sample using the hot soxhlet extraction with n-hexane, ethyl acetate, acetone, methanol and water sequentially in order of their polarity index. The extracts were tested against some clinical isolates of *Eschericia coli*, *Staphylococcus aureus*, *Aspergillus niger* and *Candida albicans*. The results showed that the water stem bark extracts of *Khaya senegalensis* W(S) B showed the highest activity with 20.00ug/ml, while the Acetone roots of *Anogeissus leiocarpus* extracts

A(R)B had the least activity with 0.00mg/ml. In case of the antibacterial activity while in the case of the antifungal activity the water stem extracts of *Khaya senegalensis* had the highest activity with 15.00ug/ml, while the Acetone roots of *Anogeissus leiocarpus*, methanol roots of extracts of *Khaya senegalensis* A(R)B, M(R)A and W(R)A had the least activity of 0.00, 0.00, and 0.00 ug/ml antifungal activity. Streptomycin Sulphate Powder was used as the standard for the antibacterial activity. while Fulcin was used as the standard for the antifungal activity. The antimicrobial activity of the roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus* justifies the claims by the traditional healers that the roots and stem barks of the plants are used to cure different ailments.

**KEY WORDS:** *Khaya senegalensis*, Streptomycin, Polarity, *Anogeissus* and Fulcin.

## INTRODUCTION

The use of plants as medicine to cure illness or diseases and to lubricate the wheel of social interaction at interpersonal- level is a behavior that predicts civilization. It is found in every society irrespective of its level of development and sophistication (Odugbemi,2006). Medicinal plants contains numerous biologically active compound such as nutrient and phytochemicals which have physiological actions on the human body (Edeoga *et al.*, 2005).

Mathew (1996) observed that 70% to 80% of the populace in Nigeria rely on plants for their primary health care needs. Today phytochemists and pharmaceutical companies depends on those medicinal plants. The potential of the higher plants as a sources of new drugs is still largely unexpressed as among the estimated 250,000 – 500,000 plants species, only a small percentage has been investigated for their phytochemical constituents and the fractions subjected to biological or pharmacological screening (Makes and Satish, 2008).

The success story of chemotherapy can only lie in one continuous search for new drugs to counter the challenges posed by resistant strain of microorganisms (Doughari *et al.*, 2008).

Studies have confirmed that extracts from *Khaya senegalensis* and *Anogeissus leiocarpus* as well as their isolated compound possess the diverse biological activities including anti-inflammatory, antitumor, antimicrobial, antioxidant, anti-ulcer to protective with alkaloid and flavones as the major active compound .

The juice obtained from the roots is applied typically for the treatment of skin infectious. In Africa stem bark extracts of *Khaya senegalensis* and *Anogeissus leiocarpus* are used to treat leprous infections, gonorrhea, syphilis and sores (Burkill,2000). In Brazil the roots extracts are used as antipyretic and anti-inflammator (Moreira, 2010). In Nigeria, the juice from both the

roots and stem barks are used for the treatment of diabetes I and II and also for the Hepatitis (Sofowora, 1993).

The specific objectives of the study includes extracting and isolating the active ingredient from both the roots and stem barks, to confirm or disprove the efficacy of the plants by evaluating their antibacterial and antifungal activity.

## **MATERIALS AND METHODS**

### **Collection of the Sample**

*Khaya senegalensis* and *Anogeissus leicocarpus* roots and stem barks were collected from their natural habitat of the coastal plain of G.R.A Sir Khasim a metropolis of Bauchi in the month of February, 2019. The samples were air dried for about two (2) weeks under shade and milled in to fine powder using Wiley milling machine. 100g of the pulverized sample of the extracts were weighed and extracted using the hot solvent extraction with n-hexane, ethyl acetate, acetone, methanol and distilled water sequentially in order of their polarity index. The extracts were stored in a desiccator for the use.

### **MICROORGANISMS**

The microorganisms for the test includes *Eschericia coli*, *staphylococcus aureus*, *Aspergillus niger* and *Candida albicans*. They were obtained from the reference isolates/organisms of Medical Microbiology unit of Abubaka Tafawa Balewa University Teaching Hospital, Bauchi (ATBUTH). The choice of these pure isolates/microorganisms was based on their implications in human diseases such as Typhoid fever, Pneumonia, Urinary infections, diabetes, hepatitis etc.

## ANTIMICROBIAL TEST

The antimicrobial activity was carried out on the crude extracts against the microorganisms. It was used as a guide to determine the active components of the roots and stem barks of *Khaya senegalensis* and *Anogeissus leicarpus*. The procedures of water birth (1978) and Perez *et al.* (1990) were employed with small modifications. The methods involved the preparation of the culture medium and inoculations. Aseptic technique was used to avoid contamination.

## PREPARATION OF THE MEDIA

Two media were employed for this research, Nutrient Agar (NA) for bacteria culture and Malt extract agar (MEA) for fungi culture. The media was prepared by dissolving 14g of NA in 1000cm<sup>3</sup> of distilled water while 25g of MEA was dissolved in 1000cm<sup>3</sup> of distilled water. They were sterilized at 121<sup>0</sup>C for 15 minutes in an auto clave and subsequently allowed to cool to about 45<sup>0</sup>C (temperature at which the agars remains molten) and formed in a plate (petri dish) and allowed to gel or solidified.

## STANDARDIZATION OF INOCULUMS

The four test organisms were sub cultured with nutrient both using both a wire loop (aseptically) and incubated for 24h r at 35<sup>0</sup>C for bacterial and 48h at 25<sup>0</sup>C for fungi. The growth of the microorganisms in the broths by turbidity produced was adjusted to march 0.1mc far land standard (10<sup>9</sup>cfu/mc) which was further changed to 10<sup>6</sup> cfu/ml for both the pathogens.

## APPLICATION OF THE EXTRACTS AND INNOCULATION OF THE PLATE

The Agar plates NA and MEA were inoculated by spreading a small volume of (0.04 to 0.10ml) of

the liquid inoculums (Sub cultured nutrient broth) by a means of glass root. One microbe was inoculated to each plates making a total of six (6) plates for each microbes. Six (6) wells for the acetone, methanol and water extracts and two (2) for the control (streptomycin for bacteria and fulcin for fungi). The inoculated plates were left for several hours in order to allow the extracts to diffuse into the agar. The nutrient agar and malt extract agar were incubated at 37<sup>0</sup>C for 24 hours for bacteria and 36 hours for fungi. The diameter of the zone of inhibition were determined using the vernier caliper and it was recorded.

## RESULTS

**TABLE 1:** The zone inhibition (mm) of the antibacterial activity of the crude extracts of acetone roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus*.

TEST	Con (mg/ml)	A(R)A	A(R)B	A(S)A	A(S)B	STREPTOMYCIN
<i>Staphylococcus aureus</i>	10 <sup>-1</sup>	12.00	10.00	12.00	14.00	0.00
	10 <sup>-2</sup>	11.00	7.00	10.00	12.00	0.00
	10 <sup>-3</sup>	7.00	4.00	8.00	10.00	0.00
	10 <sup>-4</sup>	9.00	2.00	6.00	8.00	0.00
<i>Eschericia coli</i>	10 <sup>-1</sup>	7.00	5.00	14.00	16.00	0.00
	10 <sup>-2</sup>	5.00	4.00	12.00	14.00	0.00
	10 <sup>-3</sup>	4.00	3.00	10.00	12.00	0.00
	10 <sup>-4</sup>	3.00	0.00	10.00	10.00	0.00

Keys: A(R)A – Acetone roots of *Khaya senegalensis*

A (R)B – Acetrone roots of *Anogeissus leiocarpus*

A (R)B – Acetone stem of *Khaya senegalensis*

A (R)B – Acetone stem of *Anogeissus leiocarpus*

SM - Streptomycin (control).

**Table 2:** Zone of inhibition (mm) of the antibacterial activity of the crude extracts of methanolic extracts of the roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus*

TEST obtained	Con (mg/ml)	M(R)A	M(R)B	M(S)A	M(S)B	STREPTOMYCIN
<i>Staphylococcus aureus</i>	$10^{-1}$	10.00	9.00	12.00	15.00	0.00
	$10^{-2}$	8.00	9.00	10.00	13.00	0.00
	$10^{-3}$	6.00	6.00	8.00	10.00	0.00
	$10^{-4}$	6.00	4.00	8.00	10.00	0.00
<i>Eschericia coli</i>	$10^{-1}$	11.00	11.00	13.00	16.00	0.00
	$10^{-2}$	10.00	9.00	11.00	14.00	0.00
	$10^{-3}$	10.00	6.00	8.00	12.00	0.00
	$10^{-4}$	8.00	4.00	6.00	8.00	0.00

M (R)A- Methanol roots of *Khaya senegalensis* M (R)B -Methanol roots of *Anogeissus leiocarpus* M (S)A- Methanol stem of *Khaya senegalensis* M (S)B - Methanol stem of *Anogeissus leiocarpus* ST- Streptomycin (control).


**Table 3:** Zone of inhibition (mm). The antibacterial activity of the crude extract of water, extracts of water roots steam of *Khaya senegalensis* and *Anogeissus leiocarpus*

TEST organisms	Con (mg/ml)	M(R)A	M(R)B	M(S)A	M(S)B	STREPTOMYCIN
<i>Staphylococcus Aureus</i>	$10^{-1}$	14.00	12.00	17.00	18.00	0.00
	$10^{-2}$	12.00	10.00	15.00	15.00	0.00
	$10^{-3}$	12.00	8.00	13.00	12.00	0.00
	$10^{-4}$	10.00	6.00	10.00	12.00	0.00
<i>Eschericia coli</i>	$10^{-1}$	11.00	10.00	18.00	20.00	0.00
	$10^{-2}$	10.00	9.00	18.00	18.00	0.00
	$10^{-3}$	8.00	6.00	16.00	16.00	0.00
	$10^{-4}$	6.00	4.00	12.00	14.00	0.00

W(R)A –Water roots of *Khaya senegalensis* W(R)B -Water roots of *Anogeissus leiocarpus*  
W(S)A- Water stem of *Khaya senegalensis* W(S)A- Water stem of *Anogeissus leiocarpus* ST – Streptomycin (control)

**Table 4:** Zone of inhibition (mm) of antifungal activity of the crude extracts of the Acetone roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus*.

TEST organisms	Con (mg/ml)	A(R)A	A(R)B	A(S)A	A(S)B	FL
<i>Candida albicans</i>	$10^{-1}$	6.00	7.00	10.00	12.00	0.00
	$10^{-2}$	2.00	5.00	8.00	10.00	0.00
	$10^{-3}$	0.00	3.00	6.00	6.00	0.00
	$10^{-4}$	4.00	2.00	4.00	4.00	0.00
<i>Aspergillus niger</i>	$10^{-1}$	4.00	5.00	8.00	9.00	0.00
	$10^{-2}$	2.00	3.00	6.00	7.00	0.00
	$10^{-3}$	0.00	2.00	4.00	5.00	0.00
	$10^{-4}$	0.00	0.00	2.00	2.00	0.00



PLATES 5: The zone of Inhibition of A(S)B and A(S)A on *Eschericia coli*

PLATES 7: The zone of Inhibition of W(S)A and W(S)B on *Candida albicans*  
PLATES 3: The zone of Inhibition of A(R)A and A(R)B on *Eschericia coli*


PLATES 1: The zone of Inhibition of M(R)A and M(R)B on *Escherica coli*

PLATES 6: The zone of Inhibition of Antibiotic (streptomycin) used as the positive control and water as the negative control  
PLATES 4: The zone of Inhibition of A(S)B and A(S)A on *Candida albicans*

PLATES 2: The zone of Inhibition of A(R)A and A(R)B on *Candida albicans*

PLATES 8: The zone of Inhibition of W(R)A and W(R)B on *Candida albicans*





PLATES 9: The zone of Inhibition of M(R)A and M(R)B on *Candida albicans*

PLATES 10: The zone of Inhibition of W(R)A and W(R)B on *Escherichia coli*





PLATES 11: The zone of Inhibition of W(S)A and W(S)B on *Escherichia coli*

IJSER

PLATES 12: The zone of Inhibition of M(S)A and M(S)B on *Escherichia coli*

## DISCUSSION

Antimicrobial activity of the roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus* was carried out using the clinical isolate of *staphylococcus aureus*, *Eschericia coli*, *Candida albicans* and *Aspergillus niger*.

Monica (1984) pointed out that the active antimicrobial compound diffuses from the disc into the medium and the organism are sensitive to the active antimicrobial compound are inhibited at distance from the disc. The antimicrobial activity of the roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus* was against the growth of the microorganisms (microbes) were observed visually and measured. Adewumi and Sofowora (1980) pointed out that the roots and extracts of *Khaya senegalensis* possess antimicrobial actions at different concentration depending on the bacterial species. The measured zone of inhibition of the pathogens by the crude extracts are summarized in (Table 1 – 6) and (plates 1 – 11).

The antibacterial and antifungal assays were performed for acetone, methanol and water fractions of the roots and stem barks of *Khaya senegalensis* and *Anogeissus leiocarpus* respectively against the chemical isolates of *staphylococcus aurens*, *Escherice coli*, *Aspergillus niger* and *candida albicans*.

Table 1 shows the result resists of the zone inhibition of the extracts on the clinical isolates. This proves that extracts were active at appropriate concentration against the microbes with the increase in the zone of inhibition tends to reduces as in the case of acetone extracts (Table 1).

The invitro antimicrobial activities results obtained from the roots and stem barks of *Khaya senegalensis* exhibited antibacterial activities (Table 1 – 3) with the exceptions of the C.

*abicans* and *A. niger* which had the least effects with the decrease in the concentration (Table IV).

Methanol and water extracts of the roots and stem of *Khaya senegalensis* and *Anogeissus leiocarpus* exhibited a significant antibacterial activities on the *S. aureus* and *E. coli* (Table II and III), while showed the least anti-fungal activities (Table V and VI).

The dimension of the zone of inhibition obtained for all the extracts all obtained of the (Plates 1 – 11).

Abad *et al.* (2009) Have listed *saponins Terpenoids, Flavonoids, phenolics* and steroidal compounds, are among the phytochemicals with antimicrobial effects and there phyto compounds are present in all the extracts.

The results showed increase in the antibacterial activity with increase in polarity of the solvents used in the extraction. They could mean that the concentrations of the phytochemical are higher in the polar solvent extract than the least (polar extracts). It is also noted that phyto-compounds are not presents or they are less in the acetone fractions (Table IV). It is known that compound of plants origin do not have the same antimicrobial and antifungal effects on individual organisms while some could be highly antibacterial some could be highly antifungal.

## REFERENCES

- Adewumi, A.O; Sofonora, S.A. 1980 Preliminary screening of some plants extracts for molluscidal activity plantamedica.
- Arif. T. 2009. Natural products Antifungal agents from plants. A journal of Asian Natural Products Research. 2 (7): 621 – 638.
- Conserva, L.M; Ferreira, J.C. 2012 Borrreira and spermacoce species (*Rubia ccae*): A review of their ethnomedicinal properties, chemical constituents and biological activities. Phcog., 6:46-55.
- Cornell, D.S.; Johnston, M.C 1970 Manual of the vascular plants of Texas. Texas Research foundation. Renner, TX2, 881P.
- Edeoga H.O, Okwo D.E, Mbaebie B.O (2005). Phytochemical Constituents of some Nigerian Medicinal plants. Afr. J. Biotechnology 4:685-888.
- Monica, C. 2000. District Laboratory practice in tropical countries (part 2) UK Cambridge University press, Pp. 32-34.
- Odugbemi, T.O.; Odunayo, R.; Akinsulire, E.A.P 2007 Medicinal plants useful for malar therapy in okeigbo, Ondo state and South Western Nigeria Afr. J. Erad, complementary Alternative medicine 4(2): 191-198.
- Perez, L.; panli, M.; Bezique, P. 1990. An Antibiotic assaup by the Agar well diffusion method acta Biological medicine Experimentalis, Pp 15, 113 – 115.
- Shukla S., Mehta a., Vyas. S.P and Bajpai V. (2011). Preliminary phytochemical and Antifungal screening of various organic extracts of caesalpiwa bonducella and Romania Biotechnological letters. 16 (16): 63 – 69.
- s(2008) Medicinal plants and traditional medicine in Africa. Ibadan spectrum Books limited. Pp 171 – 172.
- Sofowora. A 1993. Medicinal plants and Traditional Medicines Africa, Spectrum Books Ibadan.