

SPECTROSCOPIC STUDY OF VALINE MOLECULE

Santosh Kumar¹, Anup Kumar²

¹Dept. of applied science IEC college of engineering & tech. Gr. noida, India

²IIMT College Gr.Noida, India

¹bhu.santosh71@gmail.com

ABSTRACT-

Valin is a type of amino acid which is essential for growth of the living organism. I have monitored the IR, Raman & electronic absorption spectra of the valine molecule in solid form. The IR spectra has been observed in the range 4000 -400 cm^{-1} in the solid as well as liquid form. Similarly Raman spectra of the same molecule has also been monitored under the same range. The electronic absorption spectra has been monitored in the range 400-200 nm. In this paper we have made the several assignment of observed peaks. There are some new peaks has been found, which were not reported by earlier workers.

INTRODUCTION:

Valine molecule is a type of amino acids. It contains 19 atoms and has 51 modes of vibrations of this molecules are infrared as well as Raman active. Stepanion et al (1) studied the conformational behavior of non ionized α -valine. The structure and relative energies of the nine lowest energy valine conformers were calculated at the DFT/ B3LYP/6-31+G and MP2/6-31 + +G** and MP2/6-31++G** levels of theory. These authors (1) also recorded and analyzed the matrix isolation IR spectrum of valine which allowed a definite identification of bands corresponding to three lowest energy valine conformers I, II and III. By comparing the calculated and the observed intensity of bands for different conformers, they estimated the conformational conformation of valine in Ar matrix. They concluded that almost 94% valine is in conformer I while 5% of these are in conformer II and the remaining portion is in III conformation a number of spectroscopic studies on valine and their C-deutrated analogues [2,11], L-valine nitrate [3,12], tri L-valine selenate [4,9], have been reported in the literature. Raman optical activity of L-valine has been reported by Gargo et al [5,6&7]. Stepnain et al [1] have assigned a high frequency band at 3561 cm^{-1} due to OH stretching vibration for conformer I and at 3183 cm^{-1} due to the same OH stretch for conformer II. They also observe a band at 3422 cm^{-1} and assigned to asymmetric NH_2 stretching vibration. Another region which provides information on the valine conformational composition is the C=O stretching vibration [1800-1700 cm^{-1}]. An intense

band in this region is seen at 1761 cm^{-1} and is attributed to the lower energy conformer I in the region below 1500 cm^{-1} most of the experimental bands are assigned to the lowest energy conformer I with some characteristic bands of conformer II. The structural and electronic properties of L valine have also been studied by tulip et al [8,9&10]. They have also shown that amino and carboxy functional groups are the dominant factors in determining band structure [13,14]. In the present paper I have reported spectra of valine molecule. Large number of bands are observed in the structure some of the peaks observed here are new and not reported by earlier workers. The IR, Raman, and electronic absorption spectrum of the valine molecule are shown in Fig 1, 2 & 3 respectively.

Result and discussion-

A broad band appears in the IR spectrum of valine at 2927 cm^{-1} and is due to C-H stretch in CH_2 group. There is no corresponding peak is seen in Raman spectrum. Similarly in the Raman spectrum four peaks are seen at 2880, 2865, 2835, 2810 cm^{-1} and are attributed to CH_2 stretch. The peaks at 2835 and 2880 cm^{-1} are new a peak is observed in Raman spectrum at 2585 cm^{-1} . This is perhaps due to combination (1152+1448) cm^{-1} . A broad band is seen at 2104 cm^{-1} in IR spectrum. This is perhaps due to a combination of CH_3 bend and rocking vibration of NH_2 group (1065+1032) cm^{-1} . In Raman spectrum a broad band is seen at 1640 cm^{-1} . This is due to the asymmetric stretch of CO_2 group. The corresponding peak is absent in IR spectrum. A very intense band in the Raman spectrum appears at 1480

cm⁻¹ due to HCH bending. A band in IR spectrum at 1423 cm⁻¹ and its corresponding band at 1448 cm⁻¹ in Raman

S. N O.	I.R. frequency (cm ⁻¹)	Raman frequency (cm ⁻¹)	Assignment
1	2927	2880	Asym CH ₂ Stretch
2	-----	2865	Asym CH ₂ stretch
3	-----	2835	Sym CH ₂ Stretch
4	-----	2810	Sym CH ₂ Stretch
5	-----	2710	CH Stretch
6	-----	2585	Combi (1152+1448)
7	2104	-----	Combi (1065+1032)
8	-----	1640	CO ₂ Asym Stretch
9	-----	1480	HCH Bend
10	1423	1448	HCH Bend
11	1394	1400	OH Bend + CH Bend
12	1352	-----	CH ₃ Bend
13	1329	1360	CC Stretch, OH Bend
14	1271	1288	CH Bend
15	1177	1200	CH Bend
16	1140	1152	CN Stretch, CC Stretch
17	1065	1080	CH ₃ Bend
18	1032	1048	CH ₃ Bend
19	-----	968	CC Stretch, CH Bend
20	949	928	CN Stretch
21	900	872	NH ₂ Bend
22	823	840	OH Torsion, NH ₂
23	775	-----	Bend
24	715	710	CC Stretch, C=O Bend
25	665	633	CC Stretch, C=O Bend
26	543	----	C=O Bend, CC Bend
27	472	-----	CC Bend
28	428	435	CO Bend
			NCC Bend, CC Torsion

spectrum is due to HCH bending. A medium intense band in Raman spectrum appears at 1400 cm⁻¹ and inn IR at 1394 cm⁻¹. This is attributed to a combination of OH bend and CH bend vibrations. A weak band at 1352 cm⁻¹ in IR spectrum and at 1360 cm⁻¹ in the Raman spectrum appears due to CH₃ bending. A medium intense band appears at 1271 cm⁻¹ in IR and at 1288 cm⁻¹ in Raman and due to twisting of the CH₂ group. The CH bending vibration causes peaks in IR and Raman spectrum at 1177 cm⁻¹ and at 1200 cm⁻¹ respectively. A medium intense band appears in both the spectra at 1140 cm⁻¹ (IR) and 1152 cm⁻¹ (Raman) due to CN stretch. Similarly the peaks observed at 1065 cm⁻¹ in IR and at 1080 cm⁻¹ in Raman spectrum is due to CH₃ bending vibration. Apeak at 1032 cm⁻¹ in IR and at 1048 cm⁻¹ in the Raman spectrum peaks are seen due to CH₃bending. At 968 cm⁻¹ an intense band is seen in the Raman spectrum due to NH₂wagging. The corresponding band in the IR appears at 949 cm⁻¹. At 928 cm⁻¹ there is a band in Raman spectrum due to CN stretch .a medium intense band at 872 cm⁻¹ appears in Raman spectrum due to NH₂ bend and corresponding IR band is at 900 cm⁻¹. Due to OH torsion a peak appears at 840 cm⁻¹ in Raman spectrum and at 823 cm⁻¹ in IR spectrum. An intense band is seen at 760 cm⁻¹ in the Raman spectrum with the corresponding band in IR at 775 cm⁻¹ and is assigned to CC stretch. An intense band is observed in IR spectrum at 665 cm⁻¹ and in Raman at 633 cm⁻¹. These bands are due to C=O bending. A much intense bend is observed in IR spectrum at 543 cm⁻¹ and it may be due to CC bending. The corresponding band is not seen in Raman spectrum .in Raman spectrum as well as in the IR spectrum intense band at 435 cm⁻¹ and at 428 cm⁻¹ respectively is seen and is assigned as CC torsion.

TABLE 1-ACOMPARISON OF IR AND RAMAN FREQUENCIES OF VALINE

Asy=asymmetric, sym=symmetric, combi=combination, bend=bending

Electronic absorption spectrum of valine –

Electronic absorption spectrum of valine has been recorded in region of 200-400 nm. .01 gm of valine molecule was dissolved in 2 ml of triply distilled water. In the spectrum of this solution a peak appears at 212 nm. On further dilution (same amount of valine dissolved in 4 ml of water) a new peak appears at 204 nm .The pH of this solution is 7. On changing the pH of solution from 7 to 2 the peak remains unchanged (i.e. 204 nm). However at 10 pH a peak appears at 220 nm.




Fig-3. Electronic Absorption Spectra of Valine Molecule




Fig-1. IIR Spectra of Valine Molecule




Fig-2 Raman Spectra of Valine molecule

References:-

- [1]S.G.Stepanian,I.D.Reva,E.D.Radchenko,andI.Adamowicz ,J.Phys.chem.A103,4404,1999
- [2]S.A.Overman,G.J.ThomasJr.,Biochemistry 38,4018(1999).
- [3]B.J.M.Rajkumar,V.Ramkrishan,J.Raman Spectrosc.31,1107(2000).
- [4] I.Nemec, Z. Micka, J. mol. struct. 563, 289 (2001).
- [5] A. R. Gargo, L. D. Barron, L. Hecht, J. Raman Spectroscopy,24,91(1993)
- [6] P. R. Tulip Physical Review B71, 195117 (2005).
- [7] A.G.Caszar,J. phys.Chem. 100, 3541 (1996).
- [8] G.R.Desiraju and T.Steiner, The weak Hydrogen Bond in structural chemistry and Biology (Oxford University Press, oxford (1999).
- [9]Barone V., Carnimeo I., Scalmani G., J. Chem. Theory Comput., 2013, Vol. 9, pp. 2052-2071.
- [10]Sun J., Niehues G., Forbert H., Decka D., Schwaab G., Marx D., Havenith M., J. Am. Chem. Soc., 2014, Vol.136, pp. 5031-5038.
- [11]Debiec K.T., Gronenborn A.M., ChongL.T., J. Phys. Chem. B, 2014, Vol. 118, pp.6561-6569.
- [12]Horn A.H.C., J. Mol. Model., 2014, Vol. 20, pp. 2478-2592.

[13]Amninder S. V. , Timothy S.G., ScottA.W., Allan M.T.,
William S.P., Frontiers in Physics, 2015, Vol. 3, pp. 1-15.

[14]Bastida A., Zuniga J., Requena A., MiguelB., Candela
M.E., Soler M.A., J. Phys. Chem. B, 2016, Vol.120, pp. 348-
357.

IJSER