

Laparoscopic Cholecystectomy in Day Surgery: Feasibility and Outcomes in Taiz-Yemen

Abdulkafi Shamsan, MD¹, Tark Noman, MD, FRCS², Khulood Shukri, MS³, Maha Hizam, MD¹, Ismail Al-Shameri, MS^{3,*}

¹Department of General Surgery, AlSafwa Hospital, Faculty of Medicine, Taiz University, Yemen.

²Department of General, and cardiovascular surgery, Faculty of Medicine, Sanaa University, -Yemen

³Department of General Surgery, Authority of Althawra Hospital, Taiz University Faculty of medicine, Taiz, Yemen

BACKGROUND: Major surgery performed as a day surgery procedure is not uncommon. The aim of this study is to evaluate the feasibility of day surgery procedures in laparoscopic cholecystectomy (LC) in Taiz Yemen. **METHODS:** Between January 2017 to May 2020, in our hospital 203 patients were selected for day surgery LC. The indication for surgery was symptomatic cholelithiasis confirmed by ultrasonography. All patients were informed about the same-day discharge policy and received the postoperative instruction form on discharge. Retrospective review of Preoperative work-up included history taking and physical examination in addition to standard laboratory and radiological tests. Operative time, hospital stay, and complications were recorded. **RESULT:** 203 laparoscopic cholecystectomies were performed, the median age was 38 (IQR, 20–50) years, mean postoperative stay was 1.2 ± 0.57 days. Fifteen (7.4%) patients were discharged home at the same day. Three (1.5%) patients were re-admitted; 149 (71.8%) patients might be discharged home at same day. **CONCLUSION:** laparoscopic cholecystectomies may be done as a day surgery procedure with optimal patient satisfaction and without complications.

Keywords: Cholecystectomy, day surgery, laparoscopic, Taiz Yemen

*Corresponding author: Ismail Al-Shameri, MS, Cardiovascular center Taiz, Faculty of medicine, Taiz University, Taiz-Yemen
e-mail: Ismailamer17@gmail.com; Tel: 00967-772228396

1 INTRODUCTION

The gallbladder (GB) diseases have undergone major changes in its surgical management during the last three decades.^{1,2} Starting from open cholecystectomy, which in 1882 the first cholecystectomy is performed.^{3,4} Then a diverse technique for cholecystectomy was instituted, propagated by surgeons in France and the United States, also described briefly that it involves the use of a laparoscope and instrument insertion through trocars, thereby avoiding the classic incision.⁵ Based on the initial experiences of the surgeons with laparoscopic cholecystectomy (LC), they have become increasingly confident of discharging patients of all ages early, which saves considerable monetary and human resources.⁶

As LC is minimally invasive surgery, it facilitates faster patient recovery with minimal nursing assistance, so it is ideal to be listed as day-case procedures.⁶ Although there is lack of data

associated with day care laparoscopic cholecystectomy (DCLC) from developing countries,^{7,8} a recent review performed in a developing country reported that the choice of admitting patients overnight was mostly due to patients' preference, more than any clinical reason.⁷

Review of outcomes for LC reconfirms the established principle that LC is safe and may be performed with minimum morbidity

pooled prevalence range (1.6–5.3%) and mortality (0.08–0.14%).⁹

The aim of this is to identify clinical and surgical determinants if patient need to be kept either as 24-hour observation or same day discharge home post elective laparoscopic cholecystectomy.

Method and patients

This study is retrospective, cross section descriptive,

hospital-based study. The sample size of the study was consisting of all patient who underwent LC and patients at least with one exclusion criteria were excluded. The study period between 1st January 2017 to May 2020 in Taiz city.

Inclusion criteria

All patients of all ages and both gender who admitted to the hospital to carry out elective laparoscopic cholecystectomy.

Exclusion criteria

Patients who underwent open cholecystectomy, LC associated with other surgical procedures, as (hernioplasty and appendectomy), etc., and emergency interventions. Patients with common bile duct stones or dilatation, cholangitis, pancreatitis, significant liver function test (LFT) biochemical abnormality, and jaundice were excluded. Patients whose files that were not full filled with complete information which is needed for the study.

Data collection:

This study reviewed patients’ medical records systemically and were divided into two groups:

- a) Group with hospital stay <24 hours.
- b) Group with hospital stay ≥24 hours.

Data was collected from the hospital files using questionnaire, which was organized to five distinct sections, and were heading indicating its content; as follows:

1st Section (patient demographics): name, age, sex, address, and telephone number.

2nd Section (comorbidity variables): hypertension (HTN), diabetes mellitus (DM), ischemic heart disease (IHD). previous history of abdominal operation.

3rd Section (Pre-operative variables): information about, gallbladder stones(GBS), sludge, pericholecystic fluid, GB wall thickening, common bile duct diameter obtained from USG scan reports, **(indication of surgery)** as symptomatic GB stone, chronic cholecystitis, others if viable.

4th Section (operative variables) will be classified as (findings and complications):

- **Findings** as acute inflammation, presence of adhesion, GBS, distended GB, empyema, mucocele, and other (polyp, gangrenous GB.)
- **Complications** as incidental perforation of GB, injury of bowel by trocar, bile duct injury and abdominal drain placement, and cases that required conversion to open.

5th Section: postoperative variables and outcome will be derived from postoperative charts, will include post-operative nausea and vomiting (PONV), complications as postoperative hemorrhage (clinical signs of active hemorrhage such as bloody output from drain), bile leak (directed by drain output), and length of stay (LOS) in hospital, and patient who readmitted.

Ethics:

Data collection and research procedures in this study weren’t likely to cause any physical or emotional harm on the population under study. All data were kept private and used in the study only. Results were used only for scientific goals didn’t appear as part of this project. This protocol was submitted to ethics committee by following the applicable protocol.

Statistical analysis:

Data were processed by SPSS 25 statistical program. Quantitative variables like age were presented by calculating means ± SD. Qualitative variables like gender and LC findings were presented by calculating frequencies and percentages. The statistical significance of differences between categorical variables were calculated by the chi-square test. P-value of <0.05 was considered as statistically significant.

Result

During the period of the study, 209 patients underwent LC, out of which six patients didn’t fulfilled the inclusion criteria and two of them associated with hernia repair operation were excluded. Of the 203 patients, the median age was 38 (IQR, 20–50) years, the highest percentage 196 (96.6%) patients aged ≤ 65 years. Female 186 (91.6 %) and male 17 (8.4%) with female-to-male ratio of (11:1). 177 (87.2%) patients had no associated comorbidities.

However, hypertension was found among 10 (4.9%) patients and DM were 8 (3.9%) patients that were the commonest associated medical conditions. The findings of USG, cholelithiasis were the most common findings in 172 (84.7%) patients, in addition to 27 (13.3%) patients; which accompanied GB wall thickening more than 3 mm that show presence of acute inflammation. Two (1.0%) patients with GB wall thickening more than 3 mm one (0.5%) patient associated with polyp and one (0.5%) patient with sludge, two (1.0%) patients of GB polypoid lesions.

[Table-1] Based on medical history and abdominal imaging, the most frequency were 168 (82.8%) patients were treated for symptomatic GBS /chronic cholecystitis.

Table-1 Indications for cholecystectomy according medical history and abdominal imaging tests

Indications for surgery	N	(%)
Chronic	1	82.8

cholecystitis	68		
Acute	3		15.8
cholecystitis	2		
Gallbladder	2		1.0
polyps			
A calculous	1		0.5
cholecystitis		2	100.
Total	03	0	

[Table-2] Intraoperative complications were 6 (3%) patients; one (0.5%) patient had significant hemorrhage from cystic artery and difficulty to control it that needed to be converted to open procedure, two (1.0%) patients had stone and GB spillage during extraction lead to drain insertion and incidental perforation of GB had occurred in other 3

Table-2 Intraoperative complications among patients who underwent laparoscopic cholecystectomy

Complications	N	(%)
Significant hemorrhage	1	0.5
Incidental perforation of GB	3	1.5
stone, GB spillage	2	1.0
Total	6	3
Converted to open	8	3.9

Percentages are calculated from total number of patients (n=203)

(1.5%) patients that drain insertion is needed to be done. Eight (3.9%) patients were converted to open surgery due to difficult procedures; as the presence of significant inflammation and adhesions compromising the surgical operative field in 3 (1.5%) patients, one (0.5%) patient due to difficulty to identify cystic duct, other patient with impaction of the stone in the distal part of the cystic duct, and patient with thick wall of the GB and mucocele, and gangrenous GB in one (0.5%) patient, added to presence of adhesions in both.

[Table-3] shows post-operative variables which may contribute for Overnight stay (ONS), 21 patients complained of pain as noticed by administration of analgesic from drug list. Five patients experienced PONV as it documented at follow up note, one

patient had bile leak from accessory duct that re-admitted in the second day post discharge, other 4 (2.0%) patients have experienced different variables as hypoxia in one patient, dizziness in two patients, and asthma in one patient.

Table-3 Post-operative among patients who underwent laparoscopic cholecystectomy

Post-operative Complications	N	(%)
Bile leak	1	0.5
Pain	21	10.3
PONV	5	2.5
Other	4	2.0
Total	31	15.3

Re-admission rate was noticed with 3 (1.5%) patients, which were due to non-specific abdominal pain, sub-hepatic fluid collection which managed conservatively, and acute pancreatitis for each patient.

As all patients were admitted as inpatient, mean postoperative stay was (1.2 ± 0.57) days that was illustrated on [table-4]. Fifteen (7.4%) patients have been discharged on the same day of surgery without post-operative overnight admission. 23 (11.3%) patients of 32 (15.7%) patients who have done LC for acute cholecystitis were discharged within 24 hrs. with overnight stay and 2 (1.0%) patients were discharged at the same day according to their desire, thirteen (6.4%) patient who had LC for chronic cholecystitis have been discharged at same day.

The relation is not significant P-value (P=0.097) patients were admitted for two nights. The relation is not significant P-value (P=0.083). Re-admission rate was noticed with 3 (1.5%) patients, which were due to non-specific abdominal pain, sub-hepatic fluid collection which managed conservatively, and acute pancreatitis for each patient.

As all patients were admitted as inpatient, mean postoperative stay was (1.2 ± 0.57) days that was illustrated on [table-4]. Fifteen (7.4%) patients have been discharged on the same day of surgery without post-operative overnight admission. 23 (11.3%) patients of 32 (15.7%) patients who have done LC for acute cholecystitis were discharged

within 24 hrs. with overnight stay and 2 (1.0%) patients were discharged at the same day according to their desire, thirteen (6.4%) patient who had LC for chronic cholecystitis have been discharged at same day.

The relation is not significant P-value (P=0.097) patients were admitted for two nights. The relation is

Table- 4 Length of stay post laparoscopic cholecystectomy

Length of stay	N	%
<24 hr.	15	7.4
24 hr.	167	82.3
2 days	21	10.3
Total	203	100

Mean post-operative stay 1.2 ± 0.57 days.

Re-admissions 3 1.5

Note: 24 hr. (one-night stay), 2 Days (two-night's stay).
not significant P-value (P=0.083).

As drain had relation with length of stay (LOS) it was the most cause of inpatient, of 25 (12.3%) patients who had a drain, 11 (5.4%) patients had been admitted for two days, 14 (6.9%) patients had been admitted for one day (one-night stay). This is significant statistically (P = 0.000).

As the operation have not been planned as day case surgery, and a large different between the frequency between the two groups; so, it would be bias, and we couldn't identify with the residence of the patients is a factor for overnight stay or not.

Discussion:

Outpatient cholecystectomy was first reported by Reddick and Olsen in 1990.^{10,11} Day-case surgery in Great Britain and Ireland is defined as ‘the patient is admitted and discharged on the same day, with day surgery as the intended management. This is different from the term ‘23- hour stay’ used in the United States. Overnight stays are classed as inpatient stays in the UK.¹²⁻¹⁴

All patients in this study have done LC as inpatient, however mean postoperative stay is (1.2 ±

0.57) days,7.4% of patients have been discharged by their preference at the same day without overnight stay, 82.8% have been discharged within 24 h with

Table-5 Length of stay post laparoscopic cholecystectomy in association with post-operative variables

	LOS						Total	P-value	
	Same-day discharge		ONE DAY		TWO DAYS				
	N	%	N	%	N	%	N	%	
Pain	0	0.	15	6.9	3	2.0	18	8.9	0.156
PONV	0	0.	3	1.5	1	0.5	4	2.0	0.128
Drain	0	0.	12	5.9	9	4.4	22	10.	0.0300
Drain =Pain	0	0.	2	1.0	1	0.5	3	1.5	0.44
Total	0	0.	32	15.	1	7.4	47	24.	
			0	3	4		7		

Note: 1 day: discharged post one overnight stay

2 day: discharged post two nights stay

overnight stay. This result is slightly lesser than reported by other study, Mean postoperative stay was 1.8 ± 3.5 days, more than 78% of patients have been discharged within 24 hrs. from the operations and 22.3% have been discharged on the same day, usually 3–4 h after the end of the operation by Tebala et al.⁸in Italy 2017, as concluded by Seyednejad, Goecke, and Konkin in 2016 that 4 hrs. postoperative

observation time is sufficient for recognizing patients who require unplanned admissions.¹⁵

PONV is very unpleasant, and an exhausting problem for patients following LC, it could prolongs recovery time, delays patients' discharge usually be admitted to the hospital for observation and comfort leading to increased hospital cost.¹⁶⁻¹⁸ This study shows of 203 patient only five (2.5%) patients had mild PONV during the postoperative overnight stay, equal in number to other study, five (13%) patients had an overnight admission by Franco¹⁹ 2007, and six patients required admission as result of PONV by Lau and Brooks¹¹ 2001, study results in Kuala Lumpur, Malaysia 2015 have displayed Five patients complained of PONV in daycare group compared to tow patients in the overnight stay group¹⁷ Postoperative pain and PONV were two practical issues awaiting further improvement to facilitate ambulatory LC.²⁰

Control of pain was considered in the day case guideline.¹⁴ By Franco residual pain was in nine (24%) patients in¹⁹ Paris 2007, this study shows 21 (10.4%) patients complained of pain mostly at mid night, as they required added analgesic to the prescribed dose. Moreover; drain is one of overnight stay cause, when used, it reflects difficulty of the operation, it is useful only if left for several hours and this prevents patients from being discharged on the same day without overnight stay,⁸ in this study it is inserted in 25 (12.3%) patients; (6.9%) patients were discharged post one overnight stay, (5.4%) patients discharged post two overnight stay, with indication according to intraoperative finding, adhesions demonstrate the most percentage (4.9%), it is reasonable to postulate that the more difficult the dissection, the longer the duration of surgery and the higher the risk of complications, by Hakeem et al. 24.5% of patients in the group of unexpected overnight admission had drain inserted²¹ in UK.

Important outcome measurement in the day surgery situation are the hospital return and readmission rates, It has been suggested that an acceptable readmission rate should be between (1%-2%) and if return or readmission occurs in the first 24 hours there is a need to be analyzed as it would be emergent.²² The readmission rate in this study is 1.5% it was post 24 hours of discharge. By Lilimoe et al. (4.6%)²³ in USA, (3.4%) by Bal et al.²⁴ in New

Delhi 2003, Redamation frequency reflect whether the patient have met the condition which considered by day case surgery guide line so they don't need to back to the hospital.

My investigation is limited to some degree by its retrospective design and, BMI has not been considered within the prognostic factors in the present study. The presence or absence of comorbidities did not happen to be an independent prognostic factor for overnight stay, as they are medically controlled no complain associated to their comorbidity and most of their postoperative period passed smoothly

Just 15(7.4%) patients who discharged at the same day according to their preference. In conclusion the remaining 149(71.8%) patients have been stayed overnight without noticeable cause so, this study provides that majority of patients undergoing elective LC can be discharged home the day of surgery without overnight stay.

Conclusions:

Protocol for DCLC and day care units could be done at Taiz hospitals to start with safe day case surgery that would facilitate the procedure, make it ordinary to medical staff in hospitals, and facilitate prospective study of day case surgery therefor Patient should be educated about the same-day discharge. Same-day discharge should be considered if no drain was left at the end of the operation, and the patient free of post-operative symptoms. patient's preference should be considered in the future studies in our society.

Reference:

1. Hasbahceci M. Laparoscopic cholecystectomy: what has changed for the last three decades? *Clin Surg.* 2016;1(October 2016).
2. Gregory S, Saunders-kirkwood K, Cates JA, Services H. Open Cholecystectomy. 1993;218(2):129-137.
3. Ripetti V, Luffarelli P, Santoni S, Greco S. Laparoscopic cholecystectomy: do risk factors for a prolonged length of stay exist? *Updates Surg.* 2019;71(3):471-476.
4. Morgenstern L. Carl Langenbuch and the first cholecystectomy. *Surg Endosc.* 1992;6(3):113-114.
5. Bunge RG. JAMA revisited. *JAMA - J Am Med Assoc.*

- 2016;315(22):2474.
6. Carlomagno N, Tammaro V, Scotti A, Candida M, Calogero A, Santangelo ML. Is day-surgery laparoscopic cholecystectomy contraindicated in the elderly? Results from a retrospective study and literature review. *Int J Surg*. 2016;33:S103-S107.
 7. Ismail S, Ahmed A, Hoda MQ, Sohaib M, Zia-ur-Rehman. Prospective survey to study factors which could influence same-day discharge after elective laparoscopic cholecystectomy in a tertiary care hospital of a developing country. *Updates Surg*. 2016;68(4):387-393.
 8. Tebala GD, Belvedere A, Keane S, Khan AQ, Osman A. Day-case laparoscopic cholecystectomy: analysis of the factors allowing early discharge. *Updates Surg*. 2017;69(4):461-469.
 9. Pucher PH, Brunt LM, Davies N, et al. Outcome trends and safety measures after 30 years of laparoscopic cholecystectomy : a systematic review and pooled data analysis. *Surg Endosc*. 2018;0(0):0.
 10. Reddick EJ, Olsen DO. Outpatient Laparoscopic Laser Cholecystectomy. 1990;160(November):485-487.
 11. Lau H, Brooks DC. Predictive factors for unanticipated admissions after ambulatory laparoscopic cholecystectomy. *Arch Surg*. 2001;136(10):1150-1153.
 12. Anderson T, Walls M, Canelo R. Day case surgery guidelines. *Surgery*. 35(2):85-91.
 13. Ng L, Mercer-jones M. Day case surgery guidelines. *Surgery*. 2014;32(2):73-78.
 14. Marbaniang MJ. Day surgery guidelines. *Surgery*. 2019:1-6.
 15. Seyednejad N, Goecke M, Konkin DE. Timing of unplanned admission following daycare laparoscopic cholecystectomy. *Am J Surg*. 2017;214(1):89-92.
 16. Saunders CJ, Leary BF, Wolfe BM. Surgical Endoscopy. 1995;1(1):1263-1268.
 17. Salleh AAM, Affirul CA, Hairol O, et al. Original article Randomized Controlled Trial Comparing Daycare and Overnight Stay Laparoscopic Cholecystectomy. 2015;166(3).
 18. Psaila J, Agrawal S, Fountain U, et al. Day-surgery laparoscopic cholecystectomy: Factors influencing same-day discharge. *World J Surg*. 2008;32(1):76-81.
 19. Franco D. ORIGINAL Day-case laparoscopic cholecystectomy : results of 211 consecutive patients. 2007:421-424.
 20. Tsang Y, Poon C, Lee K, Leong H. Predictive Factors of Long Hospital Stay After. *Asian J Surg*. 2007;30(1):23-28.
 21. Hakeem AR, Oswald N, Franco F Di, Gergely S, Harris AM. ' True Day Case ' Laparoscopic Cholecystectomy in a High-Volume Specialist Unit and Review of Factors Contributing to Unexpected Overnight Stay. 2018;2018.
 22. Chauhan A, Mehrotra M, Bhatia PK, Baj B, Gupta AK. Day Care Laparoscopic Cholecystectomy : A Feasibility Study in a Public Health Service Hospital in a Developing Country. 2006;(August):1690-1695.
 23. Lillemoë KD, Lin JW, Eng M, et al. Laparoscopic Cholecystectomy as a " True " Outpatient Procedure : Initial Experience in 130 Consecutive Patients. 1998:44-49.
 24. Bal S, Reddy LGS, Parshad R, Guleria R, Kashyap L. Feasibility and safety of day care laparoscopic cholecystectomy in a developing country. *Postgrad Med J*. 2003;79(931):284-288.