

# Overview of Management Approaches of Acute Bronchitis in Children

Authors:

Badriah Omar AlOmari, Hajarkhalid AlShaqhaa, ArwaZiad AlRomaih

## Abstract:

This article covers a variety of problems surrounding acute bronchitis, including epidemiology and also pathophysiology, along with the management, prevention as well as treatment of disease in children. Much about acute bronchitis in children uncertain and this overview was taken on in order to help build a medical definition to serve as a guide for future studies. MEDLINE, AMED, EMBASE, CINAHL, and the Cochrane Library were searched in details (for articles published up to October,2017) for relevant articles in English language were included in the review.

Search terms as following “Acute bronchitis”, “bronchitis”, “chronic cough”, “Pediatric”, “Children”. Acute bronchitis is not very dangerous disease and every child at least once were sick.

The main point is distinguishing if it acute bronchitis or other disease, like pneumonia or asthma, cause symptoms are similar. That’s why family doctor should diagnose and if necessary or there is possibility of having other illnesses to do laboratory tests.

## Introduction:

Acute bronchitis is a professional term indicating a self-limited swelling of the big respiratory tracts of the lung that are called bronchi that is defined by coughing without pneumonia., which triggers raised development of mucus as well as various other modifications.

Coughing is one of the most usual sign bringing patients to the medical care doctor's workplace, and also acute bronchitis is normally the medical diagnosis in these patients. Acute bronchitis ought to be distinguished from various other usual medical diagnoses, such as pneumonia and also asthma, since these problems might require certain treatments not showed for bronchitis. Signs of bronchitis generally last regarding 3 weeks. It normally takes place as a result of a virus or breathing in things that aggravate the lungs such as cigarette smoke, fumes, dirt and also air contamination. The visibility or lack of tinted (e.g., eco-friendly) spit does not dependably separate in between viral as well as microbial reduced respiratory system infections. Bronchitis is an infection in your chest that is generally triggered by infections yet could likewise be triggered by microorganisms. Infections are accountable for greater than 90 percent of acute bronchitis infections. Prescription antibiotics are normally not indicated for bronchitis, and also must be utilized just if pertussis is suspected to lower transmission or if the patient goes to the raised threat of forming pneumonia (e.g., patients 65 years or older). The coughing might last for 2 or 3 weeks. You could likewise have high temperature, body pains, and also wheezing [1].

Viruses are normally thought to be the reason for acute bronchitis yet have actually been separated in a minority of patients [2]. Those separated in acute bronchitis (from the greatest to the least usual in collection) consist of influenza and as well as B viruses, parainfluenza virus, respiratory syncytial virus, rhinovirus, adenovirus, and also coronavirus. Human metapneumovirus has actually been determined as an original representative [3,4]. A current French research entailing grownups that had actually been immunized versus flu revealed a viral reason in 37% of 164 situations of acute bronchitis, which 21% were rhinovirus [2]. Therefore,

the return of certain virus differs in accordance with numerous aspects, consisting of the existence or lack of an epidemic, the period of the year, and also the flu inoculation condition of the populace.

Consistent microbial "infection" of the lower respiratory tracts could be a reason for illness, identified by a chronic coughing. Superb medical summaries of the illness that several currently describe as consistent or protracted microbial bronchitis (PBB) long preceded the acknowledgement that bacteria could own the illness. Physicians usually used terms such as chronic catarrh or chronic bronchitis. Throughout the very early years of the twentieth century, it was developed that a variety of microbial microorganisms were generally determined in the spit of people with chronic pulmonary illness, involving bronchiectasis and also chronic bronchitis [5].

**This article covers a variety of problems surrounding acute bronchitis, including epidemiology and also pathophysiology, along with the management, prevention as well as treatment of disease in children. Much about acute bronchitis in children uncertain and this overview was taken on in order to help build a medical definition to serve as a guide for future studies.**

### **Methodology:**

MEDLINE, AMED, EMBASE, CINAHL, and the Cochrane Library were searched in details (for articles published up to October,2017) for relevant articles in English language were included in the review. Search terms as following "Acute bronchitis", "bronchitis", "chronic

cough", "Pediatric", "Children". Reference lists of all retrieved articles were scanned for further relevant studies.


## Discussion:

- **Pathophysiology and Epidemiology**

Coughing is one of the most usual pediatric problem taken care of by Physical partitions, as well as it is much more typical in preschool kids compared to in older kids [6] Two from 3 kids aged in between 0 as well as 4 years see their FPs a minimum of once a year with acute respiratory infections, and also approximately three-quarters of them will certainly have coughing [7]. The majority of coughing are triggered by acute viral infections, as well as 7% to 12% of coughing result from asthma; all various other reasons are uncommon [8]. In Australia, 11.4% of kid coughing resulted from asthma, and also just 1.2% were brought on by pneumonia [9]; in Dutch basic method, just 1.9% of coughing was because of pneumonia [10]. With the exception of 0.3% as a result of whooping cough, all various other coughings were the outcome of acute viral infections. A lot of coughing in kids are triggered by uniform acute respiratory tract infections-- a coughing that does not conform to any kind of clear analysis symptoms such as croup, whooping coughing, pneumonia, or bronchiolitis [11].

Similar to grownups, kids' coughing, whether referred to as a signs and symptom of "top respiratory system infection" or "acute bronchitis," is one of the most often handled acute discussion in health care. These 2 medical diagnoses stand for a minimum of 75% of all coughing

seen. Of the various other reasons, asthma is one of the most usual; various other, possibly harmful, reasons are a lot less usual (Figure 1)[12].



GERD—gastroesophageal reflux disease.

**Figure 1.** Distribution of causes of acute cough among children in typical general practice[12].

Acute bronchitis was initially defined in the 1800s as swelling of the bronchial mucous membrane layers. Throughout the years, this swelling has actually been revealed to be the outcome of a different and also in some cases intricate chain of occasions. A noninfectious or contagious trigger results in bronchial epithelial injury, which triggers an inflammatory reaction with airway active responsiveness and mucus creation [13] Chosen causes that could start the cascade resulting in acute bronchitis are listed in (Table.1) [14], [15].

**Table 1.** Selected Triggers of Acute Bronchitis[14],[15].

Viruses:	adenovirus, coronavirus, coxsackievirus, enterovirus, influenza virus, parainfluenza virus, respiratory syncytial virus, rhinovirus
Bacteria:	Bordatella pertussis, Bordatellaparapertussis, Branhamellacatarrhalis, Haemophilusinfluenzae, Streptococcus pneumoniae, atypical bacteria

	(e.g., <i>Mycoplasma pneumoniae</i> , <i>Chlamydia pneumoniae</i> , <i>Legionella</i> species)
Yeast and fungi:	<i>Blastomyces dermatitidis</i> , <i>Candida albicans</i> , <i>Candida tropicalis</i> , <i>Coccidioides immitis</i> , <i>Cryptococcus neoformans</i> , <i>Histoplasma capsulatum</i>
Noninfectious triggers:	asthma, air pollutants, ammonia, cannabis, tobacco, trace metals, others

Acute bronchitis is generally triggered by a viral infection.<sup>9</sup> In patients below one year, respiratory syncytial virus, parainfluenza virus, and also coronavirus are one of the most usual isolates. In patients, one to 10 years old, parainfluenza virus, enterovirus, respiratory system syncytial virus, and also rhinovirus predominate. In patients older compared to 10 years, influenza virus, respiratory system syncytial virus, and also adenovirus are most regular. Parainfluenza virus, rhinovirus, and also enterovirus infections most frequently happen in the autumn. Influenza virus, respiratory system syncytial virus, as well as coronavirus infections, are most regular in the wintertime as well as springtime [14].

### Signs and Symptoms

Coughing is one of the most generally observed sign of acute bronchitis. The coughing starts within 2 days of infection in 85 percent of patients [16] A lot of patients have a coughing for less than 2 weeks; nonetheless, 26 percent are still coughing after 2 weeks, as well as a couple of coughing for 6 to 8 weeks [16] In case a patient's coughing matches this basic pattern, acute bronchitis must be highly presumed. In the earlier phases of the problem, kids might experience a drying, nonproductive coughing which goes forward later on to an abundant mucus-filled coughing.

The majority of medical professionals take into consideration coughing to be needed to the medical diagnosis of acute bronchitis, they differ in added demands. Various other symptoms and

signs might consist of spit manufacturing, dyspnea, wheezing, upper body discomfort, high temperature, hoarseness, malaise, rhonchi, as well as rales [17]. Single of these could exist in differing levels or might be lacking entirely. Sputum might be clear, white, yellow, green, and even tinted with blood. Peroxidase released by the leukocytes in spit triggers the color adjustments; for this reason, color alone ought to not be taken into consideration a sign of microbial infection [18]. Younger kids might have some throwing up or gagging with the coughing. The signs of bronchitis normally last 7 to 14 days, yet might likewise continue for 3 to 4 weeks. Rapid breathing, chest retractions, as well as hissing is a danger signs as well as moms and dads must straight most likely to medical professionals.

- **Physical Examination and Diagnosis**

Acute bronchitis must be distinguished from acute inflammation of the little air passages-- asthma or bronchiolitis-- which usually offers as dynamic coughing accompanied by hissing, tachypnea, respiratory system distress, as well as hypoxemia. It must additionally be differentiated from bronchiectasis [19] an unique sensation related to long-term dilatation of bronchi as well as chronic coughing. The medical diagnosis of chronic bronchitis is reserved for patients that have coughing and also spit manufacturing on many days of the month for a minimum of 3 months of the year throughout 2 successive years [20].

The physical exam of patients providing with signs and symptoms of acute bronchitis must concentrate on important indications, consisting of the visibility or lack of high temperature and also tachypnea, as well as pulmonary indications such as hissing, rhonchi, and also long term expiration. Proof of combination have to be missing. High temperature might exist in several individuals with acute bronchitis. High-grade or long term high temperature need to trigger factor to consider of pneumonia or flu [ 21].

There are no reliable analysis indicators or lab examinations, so the medical diagnosis of acute bronchitis is basically a medical one. One of the most essential problem to dismiss is acute pneumonia. This may be hard, as patients with pneumonia and also those with bronchitis could both have actually added lung noises and also high temperature (sadly, these indicators are particular neither delicate for pneumonia). Colds and also sinusitis could trigger effective coughing. Bronchitis patients typically present indications of bronchial blockage, such as hissing or dyspnea on effort. Unlike that in asthma, the swelling in acute bronchitis is short-term and also generally totally deals with quickly after the infection removes up [ 22].

**Physical exam:** The health examination could expose indicators of upper respiratory system infection, such as coryza, nasal blockage, and also pharyngeal hyperemia. There could additionally be proof of bronchial blockage (which could consist of extended expiratory stage) and also hissing, which might be highlighted forcibly expiry in the susceptible setting, or rhonchi. The existence of rales on physical exam need to prompt examination for pneumonia or CHF [23].

Bronchitis is generally detected primarily on the history as well as checkup of the kid. Several examinations might be gotten to dismiss various other illness, such as pneumonia or asthma. Additionally, the adhering to examinations might be bought to assist verify medical diagnosis [23]:

**PFT:** Pulmonary function screening is not suggested in patients with acute bronchitis. If done due to the fact that underlying asthma is suspected, medical professionals must realize that patients with acute bronchitis will certainly reveal light to modest bronchial blockage that removes with resolution of their infection as well as need to not be puzzled with asthma. PFTs



must be postponed till the patient has actually completely recouped from their infection if underlying asthma is believed.

**Laboratory research studies** are not required in the medical diagnosis of acute bronchitis. If various other medical diagnoses are thought (e.g., pneumonia), research laboratory researches might be chosen to validate these. Workplace spirometry and also lung function screening are not consistently utilized in the medical diagnosis of acute bronchitis.

**Chest X-rays** is an analysis test that applies unnoticeable electro-magnetic power beam of lights to make pictures of inner tissues, bones, and also body organs into movie.

Arterial blood gas is a test to examine the quantity of co<sub>2</sub> as well as oxygen in the blood.

**Pulse oximetry** is a tiny device that determines the quantity of oxygen in the blood. To get this dimension, a little sensing unit (like a Band-Aid) is taped into a finger or toe. When the device gets on, a little traffic signal can be seen in the sensing unit. The sensing unit is pain-free and also the red light does not fume.

**Sputum and nasal discharge societies.** An examination used to recognize the bacterium and also discover creating an infection.

**Lung (pulmonary function) examinations.** Diagnostic examinations that aid to determine the lungs' capability to exchange oxygen and also co<sub>2</sub> properly. The examinations are normally done with unique equipments that an individual should breathe into.

Approximately 12% of kids with coughing experience complications, [24] and although the difficulties are normally light and also quickly cured, some kids do end up being extremely ill. Otitis media is one of the most typical complication, complied with by rash, throwing up, and

also diarrhoea; just 5% of situations proceed to bronchitis or pneumonia. There is a scarceness of details concerning the anticipating worth of indicators and also signs in youngsters with coughing

- **Treatment**

Evidence-based suggestions have actually demonstrated that there are no effective drugs to either treatment or eliminate the signs and symptoms of acute coughing in kids [25], [26] As soon as the medical professional has actually with confidence excluded the rarer as well as much more severe problems for which treatment works, description is needed for the parents.

There excels proof that oral corticosteroids are an effective therapy for modest to serious crop [27], they minimize sign intensity, disease period, as well as return checkouts to the medical professional. Antihistamines need to be prevented, in many cases, since they dry up the secretions as well as could make the coughing even worse.

In most cases, antibiotic therapy is not essential to cure acute bronchitis, because the majority of the infections are triggered by infections. Also kids that have actually been coughing for more than 8 to 10 days generally do not require prescription antibiotics [28]. Therapy must consist of good hand hygiene and also an evasion of secondhand cigarette smoke. The majority of the therapy is helpful of the signs your youngster could have, as well as could consist of: anaesthetics, such as acetaminophen (for high temperature as well as pain), cough medicine, Increased liquid consumption, cool mist humidifier in the place could be valuable.

Therapy could focus on managing the coughing or stopping (antitussive treatment) or on making the coughing much more efficient (protussive treatment) [29].

Protussive treatment is illustrated when coughing need to be motivated (e.g., to clean the air passages of mucus). In randomized, double-blind, placebo-controlled researches of protussives in

patients with coughing for different reasons, just terbutaline (Brethine), amiloride (Midamor), and also hypertonic saline aerosols confirmed effective [30].

Antitussives could work therapies for acute management of serious coughing. They are frequently integrated with various other agents such as guaifenesin (an expectorant). Antitussive option is based upon the reason for the coughing. An antihistamine would certainly be made use of to deal with coughing linked with sensitive rhinitis, an antihistamine or a decongestant would certainly be chosen for coughing linked with postnasal water drainage, as well as a bronchodilator would certainly be ideal for coughing linked with asthma worsenings. Nonspecific antitussives, such as hydrocodone (e.g., in Hycodan), dextromethorphan (e.g., Delsym), codeine (e.g., in Robitussin A-C), carbetapentane (e.g., in Rynatuss), as well as benzonatate (e.g., Tessalon), merely reduce coughing [29] Picked nonspecific antitussives as well as their doses are provided in (Table 2) [31].

**Table2:** Selected Nonspecific Antitussive Agents [31]. Doses adjusted per manufacturer's instructions.

Hydromorphone-guaifenesin (e.g., Hycotuss)	5 mg per 100 mg per 5 mL (one teaspoon)*	Sedation, nausea, vomiting, respiratory depression
Dextromethorphan (e.g., Delsym)	30 mg every 12 hours	Rarely, gastrointestinal upset or sedation
Hydrocodone (e.g., in Hycodan syrup or tablets)	5 mg every 4 to 6 hours	Gastrointestinal upset, nausea, drowsiness, constipation
Codeine (e.g., in Robitussin A-C)	10 to 20 mg every 4 to 6 hours	Gastrointestinal upset, nausea, drowsiness, constipation

Carbetapentane (e.g., in Rynatuss)	60 to 120 mg every 12 hours	Drowsiness, gastrointestinal upset
Benzonatate (Tessalon)	100 to 200 mg three times daily	Hypersensitivity, gastrointestinal upset, sedation

- **Primary Prevention**

Therapy patients whose parents smoke need to terminate cigarette usage or smoke far from childrens, cause it is one of the most reliable methods to avoid acute bronchitis or make faster the healing process. Additionally, there is some proof that using vitamin A and also vitamin D can decrease the risk of acute bronchitis and also various other respiratory system infections [32], [33].

 **Conclusion:**

Acute bronchitis is not very dangerous disease and every child at least once were sick. The main point is distinguish if it acute bronchitis or other disease, like pneumonia or asthma, cause symptoms are similar. That's why family doctor should diagnose and if necessary or the there is possibility of having other illnesses to do laboratory tests.

 **Reference:**

1. Macfarlane J, Holmes W, Gard P, et al. Prospective study of the incidence, etiology and outcome of adult lower respiratory tract illness in the community. *Thorax* 2001;56:109-114.
2. Freymuth F, Vabret A, Gouarin S, et al. Épidémiologie et diagnostic des infections à virus respiratoire syncytial de l'adulte. *Rev Mal Respir* 2004;21:35-42.
3. Boivin G, Abed Y, Pelletier G, et al. Virological features and clinical manifestations associated with human metapneumovirus: a new paramyxovirus responsible for acute respiratory-tract infections in all age groups. *J Infect Dis* 2002;186:1330-1334.
4. Bastien N, Ward D, Van Caesele P, et al. Human metapneumovirus infection in the Canadian population. *J Clin Microbiol* 2003;41:4642-4646.
5. Hastings TW, Niles WL. The bacteriology of sputum in common non-tuberculous infections of the upper and lower respiratory tracts. *J Exp Med* (1911) 13:638–51.10.1084.
6. Morrell DC. *J R Coll Gen Pract* Symptom interpretation in general practice.. 1972 May; 22(118):297-309.
7. McCormick A, Fleming D, Charlton J. Morbidity statistics from general practice—fourth National Morbidity Survey, 1991–92. London, UK: HMSO, Office for National Statistics; 1995
8. Ayres JG, Noah ND, Fleming DM. Incidence of episodes of acute asthma and acute bronchitis in general practice 1976-87. *Br J Gen Pract*. 1993 Sep; 43(374):361-4.
9. Meza RA, Bridges-Webb C, Sayer GP, Miles DA, Traynor V, Neary S. The management of acute bronchitis in general practice: results from the Australian Morbidity and Treatment Survey, 1990–1991. *Aust Fam Physician*. 1994;23(8):1550–3
10. Verheij TJ, Kaptein AA, Mulder JD. Acute bronchitis: aetiology, symptoms and treatment. *Fam Pract*. 1989;6(1):66–9.
11. Davy T, Dick PT, Munk P. Self-reported prescribing of antibiotics for children with undifferentiated acute respiratory tract infections with cough. *Pediatr Infect Dis J*. 1998;17(6):457–62.
12. McCormick A, Fleming D, Charlton J. Morbidity statistics from general practice—fourth National Morbidity Survey, 1991–92. London, UK: HMSO, Office for National Statistics; 1995.
13. Treanor JJ, Hayden FG. Viral infections. In: Murray JF, ed. *Textbook of respiratory medicine*. 3d ed. Philadelphia: Saunders, 2000:929–84.
14. Hueston WJ, Mainous AG III. Acute bronchitis. *Am Fam Physician*. 1998;57:1270–6,1281–2.
15. Gonzales R, Steiner JF, Sande MA. Antibiotic prescribing for colds, upper respiratory tract infections, and bronchitis by ambulatory care physicians. *JAMA*. 1997;278:901–4.
16. Chesnutt MS, Prendergast TJ. Lung. In: Tierney LM, ed. *Current medical diagnosis & treatment*, 2002. 41st ed. New York: McGraw-Hill, 2002:269–362.
17. Mufson MA. Viral pharyngitis, laryngitis, croup and bronchitis. In: Goldman L, Bennett JC, eds. *Cecil Textbook of medicine*. 21st ed. Philadelphia: Saunders, 2000:1793–4.

18. Chodosh S. Acute bacterial exacerbations in bronchitis and asthma. *Am J Med.* 1987;82(4A):154–63.
19. Barker AF. Bronchiectasis. *N Engl J Med* 2002;346:1383-1393.
20. Brunton S, Carmichael BP, Colgan R, et al. Acute exacerbation of chronic bronchitis: a primary care consensus guideline. *Am J Manag Care* 2004;10:689-696.
21. Blinkhorn RJ Jr. Upper respiratory tract infections. In: Baum GL, ed. *Textbook of pulmonary diseases*. 6th ed. Philadelphia: Lippincott-Raven, 1998:493–502.
22. Metlay JP, Schulz R, Li YH, et al. Influence of age on symptoms at presentation in patients with community-acquired pneumonia. *Arch Intern Med* 1997;157:1453-1459.
23. Williamson HA., Jr Pulmonary function tests in acute bronchitis: evidence for reversible airways obstruction. *J Fam Practice.* 1987;25:251–6.
24. Hay AD, Wilson AD. The natural history of acute cough in children aged 0 to 4 years in primary care: a systematic review. *Br J Gen Pract.* 2002;52(478):401–9.
25. Irwin RS, Baumann MH, Bolser DC, Boulet LP, Braman SS, Brightling CE, et al. Diagnosis and management of cough executive summary: ACCP evidence-based clinical practice guidelines. *Chest.* 2006;129(1 Suppl):1S–23S.
26. Chang AB, Landau LI, Van Asperen PP, Glasgow NJ, Robertson CF, Marchant JM, et al. Cough in children: definitions and clinical evaluation. *Med J Aust.* 2006;184(8):398–403.
27. Kairys SW, Olmstead EM, O'Connor GT. Steroid treatment of laryngotracheitis: a meta-analysis of the evidence from randomised trials. *Pediatrics.* 1989;83(5):683–93.
28. Smucny J, Fahey T, Becker L, Glazier R, McIsaac W. Antibiotics for acute bronchitis. *Cochrane Database Syst Rev.* 2000;(4):CD000245.
29. Irwin RS, Boulet LP, Cloutier MM, Fuller R, Gold PM, Hoffstein V, et al. Managing cough as a defense mechanism and as a symptom. A consensus panel report of the American College of Chest Physicians. *Chest.* 1998;114(2 suppl managing):S133–81.
30. Irwin RS, Curley FJ, Bennett FM. Appropriate use of antitussives and protussives. A practical review. *Drugs.* 1993;46:80–91.
31. *Physicians' desk reference.* 56th ed. Montvale, N.J.: Medical Economics, 2002.
32. Chen H, Zhuo Q, Yuan W, et al. Vitamin A for preventing acute lower respiratory tract infections in children up to seven years of age. *Cochrane Database Syst Rev.* 2008;(1):CD006090
33. Ginde AA, Blatchford P, Breese K, et al. High-dose monthly vitamin D for prevention of acute respiratory infection in older long-term care residents: a randomized clinical trial. *J Am Geriatr Soc.* 2017;65:496-503.