

Properties of Intuitionistic Fuzzy Soft Matrices

S. Ruban Raj¹, M. Saradha²

Abstract - The concept of soft set is one of the recent topics developed for dealing with the uncertainties present in most of our real life situations. The parameterization tools of soft set theory enhance the flexibility of its applications. In this work, we give definition of intuitionistic fuzzy soft matrix, and their properties and examples. We also present the definitions of the subset hood of soft intuitionistic fuzzy matrices along with illustrative examples.

Keywords: Fuzzy Soft Set, Fuzzy soft matrix, Intuitionistic Fuzzy soft matrix, Cardinality of a intuitionistic fuzzy soft matrix, Degree of Sub matrix hood.

1. INTRODUCTION

Most of our real life problems in medical sciences, engineering, management, environment and social sciences often involve data which are not always all crisp, precise and deterministic in character because of various uncertainties typical for these problems. Such uncertainties are usually being handled with the help of the topics like probability, fuzzy sets, intuitionistic fuzzy sets, interval mathematics and rough sets etc. In this paper we introduced new concepts of soft intuitionistic fuzzy set. There has been incredible interest in the subject due to its diverse applications, ranging from engineering and computer science to social behavior studies. Fuzzy set was introduced by Lotfi A. Zadeh [1]. An Intuitionistic fuzzy set was provided by Atanassov [11]. However, Molodstov [2] has shown that each of the above topics suffers from some inherent difficulties due to inadequacy of their parameterization tools and introduced a concept called "Soft Set Theory" having parameterization tools for successfully dealing with various types of uncertainties. The absence of any restrictions on the approximate description in Soft Set Theory makes this theory very convenient and easily applicable. Fuzzy set theory proposed by Professor L.A. Zadeh[1] in 1965 is considered as a special case of the soft sets. In 2003, P.K.

Maji, R. Biswas and A.R. Roy[6] studied the theory of soft sets initiated by Molodstov [2] and developed several basic notions of Soft Set Theory. At present, researchers are contributing a lot on the extension of soft set theory. In 2005, Pei and Miao[7] and Chen et al. [5] studied and improved the findings of Maji et al [6]. In 2011, T.J. Neog and D.K. Sut[8] studied the theory of soft sets initiated by Molodstov [2] and developed several basic notions of Soft Set Theory. Recently, Cagman et al. [9] introduced soft matrix and applied it in decision making problems. In one of our earlier work [10], we proposed the idea of 'Properties of Soft Intuitionistic Fuzzy Set' in sequel to [10] defining some operations. The present paper aims to define intuitionistic fuzzy soft matrices and establish some results on them. In this paper, we define a submatrixhood of intuitionistic soft fuzzy matrices along with several examples.

1. Associate Professor, Department of Mathematics, St. Joseph College, Trichirappalli, Tamilnadu

Email:ruban946@gmail.com

2. Research Scholar, Department of Mathematics, St. Joseph College, Trichirappalli, Tamilnadu

Email:ishuharisri88@gmail.com

2. Preliminaries:

2.1. Definition [2]

Let X be an initial set and U be a set of parameters. Let $P(X)$ denotes the power set of X , and let $A \subset U$. A pair (F, A) is called a soft set over X , where F is a mapping given by $F: A \rightarrow P(X)$.

In other words, a soft set over X is a parameterized family of subsets of the universe X . For $\xi \in A$, $F(\xi)$ may be considered as the set of ξ -elements of the soft set (F, A) or as the ξ - approximate elements of the soft set. Clearly, a soft set is not a (crisp) set. To illustrate this idea, let we consider the following example.

Example:

Let us consider a soft set (F, A) which describes the “attractiveness of houses” that Mr. P is considering to purchase.

Suppose that there are six houses in the universe X under consideration,

$$X = \{h_1, h_2, h_3, h_4, h_5, h_6\}$$

and $A = \{e_1, e_2, e_3, e_4, e_5\} \subset U$

is a set of decision parameters, where e_1 stands for the parameters “expensive”, e_2 stands for the parameters “beautiful”, e_3 stands for the parameters “wooden”, e_4 stands for the parameters “cheap”, e_5 stands for the parameters “in the green surrounding”.

Suppose that

$$F(e_1) = \{h_3, h_5\}, F(e_2) = \{h_1, h_3, h_5\}, F(e_3) = \{h_1, h_4, h_5\}, F(e_4) = \{h_1, h_2\}, F(e_5) = \{h_5\}.$$

Therefore, $F(e_1)$ means “houses (expensive)”, whose functional value is the

set $\{h_2, h_5\}$. Thus, we can view the soft set (F, A) as a collection of approximations as below

$$(F, A) = \left\{ \begin{array}{l} \text{expensive houses} = \{h_2, h_5\} \\ \text{beautiful houses} = \{h_1, h_3, h_5\} \\ \text{wooden houses} = \{h_1, h_4, h_5\} \\ \text{cheap houses} = \{h_1, h_2, h_5\} \\ \text{in green surrounding houses} = \{h_5\} \end{array} \right\}$$

2.2. Definition [6]

A pair (F, A) is called a fuzzy soft set over X where $F: A \rightarrow \tilde{P}(X)$ is a mapping from A into $\tilde{P}(X)$.

Example:

Let $X = \{h_1, h_2, h_3, h_4, h_5, h_6\}$ be the set of six houses under consideration and $A = \{e_1, e_2, e_3, e_4, e_5\} \subset U$ is a set of decision parameters,

Consider the mapping $F: A \rightarrow \tilde{P}(X)$.

$$(F, A) = \{F(e_1) = \{h_2/0.9, h_5/0.7\}, F(e_2) = \{h_1/0.6, h_3/0.9, h_5/0.7\}, F(e_3) = \{h_1/0.5, h_4/0.9, h_5/0.6\}, F(e_4) = \{h_1/0.7, h_2/0.8, h_5/0.9\}, F(e_5) = \{h_5/0.8\}\}$$

is the fuzzy soft set representing the “attractiveness of the houses” which Mr. X is going to buy.

2.3. Definition [9]

Let X be a universal set & U be the set of parameters. Let $A \subseteq U$ & (F, A) be a fuzzy soft set in the fuzzy soft class (X, U) . Then fuzzy soft set (F, A) in a matrix form as $A_{m \times n} = [a_{ij}]_{m \times n}$. Or $A = [a_{ij}]$ where $i=1, 2, \dots, m$ & $j=1, 2, \dots, n$ Where

$$a_{ij} = \begin{cases} \mu_j(x_i) & \text{if } e_j \in A \\ 0 & \text{if } e_j \notin A \end{cases}$$

$\mu_j(x_i)$ represents the membership of x_i in the fuzzy set.

Example:

Let $X = \{h_1, h_2, h_3, h_4\}$ be the set of four houses under consideration and $A = \{e_1, e_2, e_3, e_4\} \subset U$ is a set of decision parameters,

The matrix representation is,

$$\begin{pmatrix} 0.8 & 0.4 & 0.3 & 0.6 \\ 0 & 1 & 0.8 & 0.9 \\ 0.7 & 0 & 0.8 & 0 \\ 0 & 0.8 & 0.9 & 0.6 \end{pmatrix}$$

2.4. Definition

Let X be an initial universe, U be the set of parameters and $A \subseteq U$. Let (F, A) be an intuitionistic fuzzy soft matrix (IFSM) over U . Let $F(X \times Y)$ denotes the family of an intuitionistic fuzzy relationship on X to Y . The set

$\tilde{I}_\alpha = \{(x, y) \in X \times Y : \tilde{I}_{\mu_\alpha}(x, y) \geq \alpha \text{ and } \tilde{I}_{\gamma_\alpha}(x, y) \leq \alpha\} \subset X \times Y$ is defined as α -cut set if $\tilde{I} \in F(X \times Y)$ for $\alpha \in [0, 1]$. Then a subset of $X \times U$ is uniquely defined by,

$$F: A \rightarrow \tilde{F}(X) \text{ and } F(x) = \{y \in X : (x, y) \in \tilde{I}_\alpha, x \in A, y \in X, \alpha \in [0, 1]\}.$$

The membership function and non membership function are written by

$$\tilde{I}_{\mu_\alpha} : X \times U \rightarrow [0, 1] \text{ and } \tilde{I}_{\gamma_\alpha} : X \times U \rightarrow [0, 1] \text{ where } \tilde{I}_{\mu_\alpha} : (x, y) \in [0, 1] \text{ and}$$

$\tilde{I}_{\gamma_\alpha} : (x, y) \in [0, 1]$ are the membership value respectively of $x \in X$ for each $y \in Y$.

If $(\mu_{ij}, \gamma_{ij}) = (\tilde{I}_{\mu_\alpha}(x, y), \tilde{I}_{\gamma_\alpha}(x, y))$, we can define a matrix, $[(\mu_{ij}, \gamma_{ij})]_{m \times n} =$

$$\begin{pmatrix} (\mu_{11}, \gamma_{11}) & (\mu_{12}, \gamma_{12}) & \dots & (\mu_{1n}, \gamma_{1n}) \\ (\mu_{21}, \gamma_{21}) & (\mu_{22}, \gamma_{22}) & \dots & (\mu_{2n}, \gamma_{2n}) \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ (\mu_{m1}, \gamma_{m1}) & (\mu_{m2}, \gamma_{m2}) & \dots & (\mu_{mn}, \gamma_{mn}) \end{pmatrix}$$

Which is called an $m \times n$ IFSM of the IFSM (F, A) over X .

Therefore, we can say that an IFSS (F, A) is uniquely characterized by the matrix $[(\mu_{ij}, \gamma_{ij})]_{m \times n}$ and both concepts are interchangeable. The set of all $m \times n$ IFS matrices over X will denoted by $\text{IFSM}_{m \times n}$.

Example.

Suppose that $X = \{h_1, h_2, h_3, h_4, h_5, h_6\}$ is the set of houses and $U = \{\text{expensive } (e_1), \text{ beautiful } (e_2), \text{ wooden } (e_3), \text{ cheap } (e_4)\}$. If $A \subseteq U = \{e_1, e_2, e_3\}$ and $\tilde{I}_\alpha = (0.6, 0.4)/(h_1, e_1) + (0.7, 0.3)/(h_1, e_2) + (0, 1)/(h_1, e_3) + (0.9, 0.1)/(h_2, e_1) + (0, 1)/(h_2, e_2) + (0.7, 0.3)/(h_2, e_3) + (0.5, 0.5)/(h_3, e_1) + (0.7, 0.3)/(h_3, e_2) + (0.9, 0.1)/(h_3, e_3) + (0.3, 0.5)/(h_4, e_1) + (0.9, 0.1)/(h_4, e_2) + (0.6, 0.4)/(h_4, e_3) + (0.7, 0.3)/(h_5, e_1) + (0.6, 0.4)/(h_5, e_2) + (0.5, 0.5)/(h_5, e_3) + (0.8, 0.1)/(h_6, e_1) + (0.5, 0.2)/(h_6, e_2) + (0.7, 0.2)/(h_6, e_3)$.

Then, a soft intuitionistic fuzzy matrix

(F, A) is a parameterized family

$\{F(e_1), F(e_2), F(e_3)\}$ of all IFS matrices over X .

Hence the IFSM $[(\mu_{ij}, \gamma_{ij})]$ is written by

$$[(\mu_{64}, \gamma_{64})]_{6 \times 4} =$$

$$\begin{pmatrix} (0.6, 0.4) & (0.7, 0.3) & (0, 1) & 0 \\ (0.9, 0.1) & (0, 1) & (0.7, 0.3) & 0 \\ (0.5, 0.5) & (0.7, 0.3) & (0.9, 0.1) & 0 \\ (0.3, 0.5) & (0.9, 0.1) & (0.6, 0.4) & 0 \\ (0.7, 0.3) & (0.6, 0.4) & (0.5, 0.5) & 0 \\ (0.8, 0.1) & (0.5, 0.2) & (0.7, 0.2) & 0 \end{pmatrix}$$

2.5. Definition

Let $(F, A) = [(\mu_{ij}, \gamma_{ij})] \in \text{IFSM}_{m \times n}$. Then (F, A) is called

- (a) A zero IFSM, denoted by $\check{0} = [(0, 0)]$, if $\mu_{ij} = 0$ and $\gamma_{ij} = 0$ for all i and j .
- (b) A μ -universal IFSM, denoted by $\check{1} = [(1, 0)]$, if $\mu_{ij} = 1$ and $\gamma_{ij} = 0$ for all i and j .
- (c) A γ -universal IFSM, denoted by $\check{1} = [(0, 1)]$, if $\mu_{ij} = 0$ and $\gamma_{ij} = 1$ for all i and j .
- (d) $A = [(\mu'_{ij}, \gamma'_{ij})]$ is a intuitionistic fuzzy soft sub matrix of $B = [(\mu''_{ij}, \gamma''_{ij})]$, denoted by $(F, A) \subset (G, B)$ if $\mu'_{ij} \leq \mu''_{ij}$ and $\gamma'_{ij} \leq \gamma''_{ij}$ for all i and j .
- (e) $(F, A) = [(\mu'_{ij}, \gamma'_{ij})]$ and $(G, B) = [(\mu''_{ij}, \gamma''_{ij})]$ are intuitionistic fuzzy soft equal matrices, denoted by $(F, A) = (G, B)$, if $\mu'_{ij} = \mu''_{ij}$ and $\gamma'_{ij} = \gamma''_{ij}$ for all i and j .

2.6. Definition

Let $(F, A) = [(\mu'_{ij}, \gamma'_{ij})]$, $(G, B) = [(\mu''_{ij}, \gamma''_{ij})] \in \text{IFSM}_{m \times n}$. Then the IFSM

$$(H, C) = [(\mu_{ij}, \gamma_{ij})]$$
 is called

(a) Union of (F, A) and (G, B) , denoted by $(F, A) \check{\cup} (G, B)$ if $\mu_{ij} = \max \{(\mu'_{ij}, \mu''_{ij})\}$ and $\gamma_{ij} = \min \{(\gamma'_{ij}, \gamma''_{ij})\}$ for all i and j .

(b) Intersection of (F, A) and (G, B) , denoted by $(F, A) \check{\cap} (G, B)$ if

$$\mu_{ij} = \min \{(\mu'_{ij}, \mu''_{ij})\} \text{ and } \gamma_{ij} = \max \{(\gamma'_{ij}, \gamma''_{ij})\} \text{ for all } i \text{ and } j.$$

(c) Complement of (F, A) is denoted by

$$(F, A)^c = [(\gamma_{ij}, \mu_{ij})]$$
 for all i and j .

Example:

Let $(F, A) =$

$$\begin{pmatrix} (0.1, 0.2) & (0.5, 0.4) & (0.3, 0.6) \\ (0.4, 0.4) & (0.2, 0.3) & (0.5, 0.1) \\ (0.5, 0.2) & (0.3, 0.4) & (0.6, 0.2) \\ (0.7, 0.2) & (0.6, 0.1) & (0.5, 0.3) \end{pmatrix}$$

and $(G, B) =$

$$\begin{pmatrix} (0.5, 0.3) & (0.1, 0.6) & (0.7, 0.1) \\ (0.8, 0.1) & (0.4, 0.3) & (0.5, 0.2) \\ (0.2, 0.5) & (0.3, 0.6) & (0.4, 0.5) \\ (0.1, 0.7) & (0.2, 0.5) & (0.5, 0.1) \end{pmatrix}$$

Then $(F, A) \check{\cup} (G, B) =$

$$\begin{pmatrix} (0.5, 0.2) & (0.5, 0.4) & (0.7, 0.1) \\ (0.8, 0.1) & (0.4, 0.3) & (0.5, 0.1) \\ (0.5, 0.2) & (0.3, 0.4) & (0.6, 0.2) \\ (0.7, 0.2) & (0.6, 0.1) & (0.5, 0.1) \end{pmatrix}$$

$$(F, A) \check{\cap} (G, B) =$$

$$\begin{pmatrix} (0.1, 0.3) & (0.1, 0.6) & (0.3, 0.6) \\ (0.4, 0.4) & (0.2, 0.3) & (0.5, 0.2) \\ (0.2, 0.5) & (0.3, 0.6) & (0.4, 0.5) \\ (0.1, 0.7) & (0.2, 0.5) & (0.5, 0.3) \end{pmatrix},$$

and

$$(F, A)^c =$$

$$\begin{pmatrix} (0.2, 0.1) & (0.4, 0.5) & (0.6, 0.3) \\ (0.4, 0.4) & (0.3, 0.2) & (0.1, 0.5) \\ (0.2, 0.5) & (0.4, 0.3) & (0.2, 0.6) \\ (0.2, 0.7) & (0.1, 0.6) & (0.3, 0.5) \end{pmatrix}$$

3.1.Cardinality of a Fuzzy Soft Matrix:

For a finite fuzzy soft matrix (F, A), the Cardinality |(F, A)| is defined as,

$$|(F, A)| = \sum_{x \in U} \mu_{ij}(x) = \sum(F, A).$$

And its relative cardinality ||(F, A)|| is,

$$||(F, A)|| = \frac{|(F, A)|}{16}$$

3.2. Set Cardinality of a Fuzzy Soft Matrix:

We define for a finite fuzzy soft matrix (F, A), the Set Cardinality |(F, A)| is defined as,

$$|(F, A)| = \sum_{e_j \in A} \exp F(e_j)$$

Example:

Let $X = \{h_1, h_2, h_3, h_4\}$ be the set of houses under consideration and $U = \{\text{expensive } (e_1), \text{ near by city } (e_2), \text{ cheap } (e_3), \text{ beautiful } (e_4)\}$ be the set of parameters. Consider the fuzzy soft matrices (F, A) and (G, B) where $A = \{e_1, e_2, e_3\} \subseteq U$. Then

$$(F, A) =$$

$$\begin{pmatrix} (0.6, 0.3) & (0.6, 0.2) & (0.5, 0.4) & (0, 0) \\ (0.9, 0.1) & (0.8, 0.1) & (0.6, 0.3) & (0, 0) \\ (0.7, 0.2) & (0.9, 0.1) & (0.5, 0.4) & (0, 0) \\ (0.5, 0.4) & (0.7, 0.3) & (0.9, 0.1) & (0, 0) \end{pmatrix}$$

$$|(F, A)| = \sum_{e_{ij} \in A} \exp F(e_{ij}) = 3.3.$$

3.3.Degree of Subset hood of Soft Intuitionistic Fuzzy Matrices

Definition.

Let X be a Universal, U be a set of parameters and let $(F, A)_{\check{I}}$ and $(G, B)_{\check{I}}$ are two soft intuitionistic fuzzy matrix of X. Then the degree of subset hood denoted by $\mathfrak{S}(A, B)_{\check{I}}$ is defined as,

$$\mathfrak{S}(A, B)_{\check{I}_{\mu}} = \frac{1}{|(F, A)_{\check{I}_{\mu}}|} \left\{ |(F, A)_{\check{I}_{\mu}}| - \sum \max\{0, (F, A)_{\check{I}_{\mu}} - (G, B)_{\check{I}_{\mu}}\} \right\}$$

where

$$|(F, A)_{\check{I}_{\mu}}| = \sum \mu_A(x)$$

$$|F, A|_{\tilde{I}_\gamma} = \sum \gamma_A(x)$$

$$|G, B|_{\tilde{I}_\mu} = \sum \mu_B(x)$$

$$|G, B|_{\tilde{I}_\gamma} = \sum \gamma_B(x)$$

and

$$\mathfrak{S}(B, A)_{\tilde{I}_\mu} = \frac{1}{|(G, B)|_{\tilde{I}_\mu}} \left\{ |(G, B)|_{\tilde{I}_\mu} - \sum \max\{0, (G, B)_{\tilde{I}_\mu} - (F, A)_{\tilde{I}_\mu}\} \right\}$$

Example:

Let $X = \{h_1, h_2, h_3, h_4\}$ be the set of houses under consideration and $U = \{\text{expensive } (e_1), \text{ near by city } (e_2), \text{ cheap } (e_3), \text{ beautiful } (e_4)\}$ be the set of parameters. Consider the fuzzy soft matrices (F, A) and (G, B) where $A, B \subseteq U$.

Then (F, A) =

$$\begin{pmatrix} (0.8, 0.1) & (0.8, 0.2) & (0, 0) & (0.6, 0.4) \\ (0.9, 0.1) & (0.7, 0.2) & (0.4, 0.5) & (0.5, 0.4) \\ (0.8, 0.1) & (0.5, 0.4) & (0.2, 0.7) & (0.6, 0.3) \\ (0.7, 0.2) & (0.6, 0.2) & (0.1, 0.7) & (0.4, 0.5) \end{pmatrix}$$

and (G, B) =

$$\begin{pmatrix} (0.8, 0.1) & (0.9, 0.1) & (0.6, 0.3) & (0.7, 0.2) \\ (0.9, 0.1) & (0.8, 0.2) & (0.8, 0.2) & (0.6, 0.3) \\ (0.9, 0.1) & (0.8, 0.2) & (0.4, 0.6) & (0.9, 0.1) \\ (0.8, 0.1) & (0.6, 0.3) & (0.8, 0.1) & (0.7, 0.2) \end{pmatrix}$$

$$|(F, A)| = \sum_{e_{ij} \in A} \exp F(e_{ij}) = 3.2$$

$$|(G, B)| = \sum_{e_{ij} \in A} \exp G(e_{ij}) = 3.5$$

$$(F, A) - (G, B) = (-0.1, 0, -0.1, -0.1)$$

$$\mathfrak{S}(A, B) = \frac{1}{3.2} \{3.2 - \sum \max\{0, (-0.1, 0, -0.1, -0.1)\}\} = 1 \text{ and}$$

$$\mathfrak{S}(B, A) =$$

$$\frac{1}{3.5} \{3.5 \sum \max\{0, (0.1, 0, 0.1, 0.1)\}\} = 0.9143 \cong 0.9.$$

4. Conclusion:

In this article, we have explained a very important consistency principle to Degree of Sub matrix hood of the fuzzy soft matrix. In this paper the basic concept of a vague soft matrix is recalled. We have introduced the concept of sub matrix hood of the soft fuzzy matrix as an extension to the fuzzy soft matrix. The basic properties on soft fuzzy matrix are also presented. The null, union, intersection, sub matrix and sub matrix hood as far as future direction are concerned. It is hoped that our findings will help enhancing this study on fuzzy soft matrices for the researchers.

5. References:

[1] Zadeh, L. A., "Fuzzy Sets", Information and Control, 8, pp. 338-353, 1965.

[2] Molodstov. D.A., "Soft Set Theory – First Result", Computers and Mathematics with Applications, Vol. 37, pp. 19-31, 1999.

[3] Ahmad, B. and Kharal, A., "Mappings on Soft Classes" (Originally Submitted on 17th oct, 2008 to Information Sciences with the title of "Mappings on Soft Sets" and was given MS#INS-D-08-1231 by EES.

[4] Ali, M. Irfan, Feng F, Liu, X., Min, W.K. Shabir, M. "On some new operations in soft set theory", *Computers and Mathematics with Applications* 57 (2009) 1547-1553.

[5] Chen, D., Tsang, E.C.C., Yeung, D.S. and Wang, X., "The Parameterization reduction of soft sets and its applications", *Computers and Mathematics with Applications* 49 (2005) 757-763.

[6] Maji, P. K., Biswas .R, and Roy, A. R., "Soft Set Theory", *Computers and Mathematics with Applications* 45, pp. 555-562, 2003.

[7] Pei, D. and Miao, D. "From Soft Sets to Information Systems", in *Proceedings of the IEEE International Conference on Granular Computing*, vol. 2, pp. 617-621, 2005.

[8] T.J. Neog and D.K. Sut, "A New Approach to the Theory of Soft Sets", in *International Journal of Computer Applications*, vol. 32, No 2, October 2011.

[9] Cagman, N. & Enginoglu, S., "Computers & Mathematics with Applications", 2010, 59 3308-3314.

[10] Ruban Raj, S.& Saradha, M., "Properties of Soft Intuitionistic Fuzzy Set" in *International Journal of Mathematics Trends & Technology*, Volume 3 Issue 6 - 2012.

[11] Chetia, B., & Das, P.K., "Journal of the Assam Academy of Mathematics, 2010,2,71-83.