

# Solid-Liquid Extraction Kinetics of Total Phenolic, Total Flavonoid, Rutin and Tannin Contents in 50% Ethanol Extract of *Cotinus coggygria*

Valentina Lubomirova Christova-Bagdassarian<sup>1</sup>, Maria Stefanova Atanassova<sup>2</sup>, Vesna K. Hristova<sup>3</sup>, M. Ayaz Ahmad<sup>4</sup>

**Abstract**— The use of a natural product with therapeutical properties has a long history. Ethanolic extract of *Cotinus coggygria* were analyzed for its phytoconstituents show that the dry leaves of *Cotinus coggygria* are rich in total phenols, total flavonoids, rutin and tannins. Solid-liquid extraction was carried out by 50% aqueous ethanol for 120 min, which gave concentration of total polyphenols and the findings of our present results are within good agreement with the other workers.

**Index Terms**— Rutin, Tannins, Total flavonoids, Total phenolics, Solid-liquid extraction, 50% aqueous ethanol extract of Bulgarian dry leaves of *Cotinus coggygria*

## 1 INTRODUCTION

Nature has been a source of medicinal agents since time immemorial. Plants play a significant role in providing primary health care. They serve as therapeutic agents as well as important raw materials for the manufacturing of traditional and modern medicines as well as in food industries. The revival of interest in plant derived drugs is mainly due to the current widespread belief that "green medicine" is safer and more dependable than the costly synthetic drugs mainly of which have adverse side [1, 2]. The medicinal value of these plants lies in some chemical constituents that produce a definite physiological action on the human body. The most important of these bioactive constituents of plants are alkaloids, tannins, flavonoids, phenolic compounds etc [2, 3].

In recent years there is an increasing interest in finding antioxidant phytochemical because they inhibit the propagation of few diseases like atherosclerosis, diabetes, arthritis, cancer, Alzheimer, ageing, neurogenerative diseases, etc [2, 4]. The interest in the investigation of active components, especially polyphenols, from natural sources (fruits, vegetables, cereals, herbs) has greatly increased in recent years. The reason for that is restricted use of synthetic antioxidants such as butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT) in foods because of their possible undesirable effects on human health [5, 6]. The free radicals are also produced by oxidation of lipids of foods is responsible for the formation of off-flavors and undesirable chemical compounds which may be harmful to health so antioxidants are used by the food industry to delay the oxidation process.

Antioxidants act by donating H-atom or by donating electrons to radical oxygenated species. Radicals obtained from antioxidants with molecular structure such as phenol, are stable species and will stop the oxidation chain reactions [2, 7]. Many studies had revealed that phenolic content in plants is mainly responsible to their antioxidant activities. Higher the total phenolic content greater is the antioxidant power [2].

*Cotinus coggygria* is one of two species constituting a minor genus of the family Anacardiaceae, viz., *Cotinus coggygria* Scop. (syn.: *Rhus cotinus* L.) itself and *Cotinus obovatus* Raf., the American smoketree. Its wide distribution extends from southern Europe, the Mediterranean, Moldova and the Caucasus, to central China and the Himalayas [8, 9]. *C. coggygria* is a common medicinal plant (well known as 'smradlika' or 'tetra') in the Bulgarian folk medicine for outer use predominantly [10, 11]. Plants of the family Anacardiaceae have a long history of use by various peoples for medicinal and other purposes. *Rhus glabra* is traditionally used in the treatment of bacterial diseases such as syphilis, gonorrhea, dysentery and gangrene, while *R. coriaria*, besides its common use as a spice consisting of ground dried fruits with salt, is also widely used as a medicinal herb, particularly for wound healing [9, 12]. In folk medicine, *Cotinus coggygria* is routinely used as an antiseptic, anti-inflammatory, antimicrobial and antihemorrhagic agent in wound-healing [12, 13], as well as for countering diarrhea, paradontosis, and gastric and duodenal ulcers [12, 14]. However, these by-products are still a good and cheap source of high-quality polyphenolic compounds which can be used in different therapeutic procedures with the purpose of free radical neutralisation in biological systems [6, 15, 16].

Extraction is a very important stage in isolation, identification and use of phenolic compounds and there is no single and standard extraction method. Solid-liquid extraction of phenolic compounds with different solvents [6, 17, 18, 19] and in some cases extraction with supercritical fluid [6, 20] are the most common by used techniques for isolation of these compounds. Many authors investigated solid-liquid extraction of natural antioxidants and their properties from grape seed [5, 6, 16]

1. National Centre of Public Health and Analyses, Department "Public Health and Health risk", 15 Akad. Ivan Ev. Geshov blvd, 1431 Sofia, Bulgaria. E-mail: v.hristova@ncpha.government.bg

2. Metallotechnica Ltd., 63 Shipchenski prohod Bul., 1574 Sofia, Bulgaria. E-mail: stefanova@myway.com

3. Faculty of Biotechnical Sciences, University of St. Clement Ohridski, Partizanska, bb 7000 Bitola, Macedonia. E-mail: v.primavera79@gmail.com

4. Physics Department, Faculty of Science, P.O. Box 741, University of Tabuk - 71491, Saudi Arabia. E-mail: mayaz.alig@gmail.com

and from other plant materials [6, 21, 22, 23] as well as methods for their identification [6, 24, 25]. However, literature data about optimisation [6, 26, 27], modelling and simulation of solid-liquid extraction process are scarce. Therefore, there is a need for mathematical modelling, as a useful engineering tool, which considerably facilitates optimisation, simulation, design and control of processes and contributes to utilization of energy, time and solvent [6].

## 2 MATERIALS AND METHODS USED

### 2.1 Plant Materials

The leaves from *Cotinus coggygia* were harvested from different regions of Bulgaria. All sample data are stated in the sampling protocol. The dried leaves were kept in a dry place until further use.

### 2.2 Sample Preparation

A dry sample of 0.5 g was weighted and phenolic and flavonoid compounds were extracted with 50 ml 50% aqueous ethanol on an ultrasonic bath for 20 min. An aliquot (2 mL) of the extracts was ultracentrifugated for 5 min at 14 000 rpm. The extract prepared in this way was used for further spectrophotometric determination of total polyphenols.

### 2.3 Sample Preparation

#### Determination of total phenolics assay

The total phenolic content of *Cotinus coggygia* were determined by using the Folin-Ciocalteu assay. An aliquot (1 mL) of extracts or standard solution of gallic acid (10, 20, 40, 60, 80, 100 and 120 mg/L) was added to 25 mL volumetric flask, containing 9 mL of distilled deionised water (dd H<sub>2</sub>O). A reagent blank using dd H<sub>2</sub>O was prepared. One milliliter of Folin-Ciocalteu's phenol reagent was added to the mixture and shaken. After 5 min, 10 mL of 7% Na<sub>2</sub>CO<sub>3</sub> solution was added to the mixture. To the solution the dd H<sub>2</sub>O was added up to volume of 25 mL and mixed. After incubation for 90 min at room temperature, the absorbance against prepared reagent blank was determined at 750 nm with an UV-Vis Spectrophotometer BOECO – Germany. All samples were analyzed in duplicates [28].

#### Determination of total flavonoids assay

The total flavonoid contents were measured by aluminum chloride colorimetric assay. An aliquot (1 mL) of extracts or standard solution of catechin (10, 20, 40, 60, 80, 100 and 120 mg/L) was added to 10 mL volumetric flask, containing 4 mL of distilled deionised water (dd H<sub>2</sub>O). To the flask was added 0.3 mL 5% NaNO<sub>2</sub>. After 5 min, 0.3 mL of 10% AlCl<sub>3</sub> was added. At 6<sup>th</sup> min, 2 mL 1 M NaOH was added and the total volume was made up to 10 mL with dd H<sub>2</sub>O. The solution was mixed well and the absorbance was measured against prepared reagent blank at 510 nm. UV-Vis Spectrophotometer BOECO – Germany. All samples were analyzed in duplicates [28].

### Rutin assay

The analyses of rutin content in *Cotinus coggygia* were performed according to The International Pharmacopoeia and AOAC method, after modified methods with 50% aqueous ethanol. Pipet 2 ml aliquots solution into 50 ml volumetric flask was added to 2 ml deionized water (dd H<sub>2</sub>O) and 5 ml ammonium molybdat. The solution was added volume (50 ml) with dd H<sub>2</sub>O and mixed. Was prepared standard solution of rutin (0.0200 g dissolved into 2 ml ethanol) was added volume (50 ml) with 50% aqueous ethanol. An aliquot (1 ml) of standard solution into 50 ml volumetric flask and dilute to volume with distilled deionized water (dd H<sub>2</sub>O). A reagent blank using dd H<sub>2</sub>O was prepared. The absorbance against prepared reagent blank was determined at 360 nm with an UV-Vis Spectrophotometer BOECO – Germany. All samples were analyzed in duplicates [29].

Calculations are based on averaging results from analyses of duplicate samples.

Calculate content (%) of rutin (R) in sample as follows:

$$R(\%) = \frac{A_{\text{sample}} \times C \times 50 \times 100}{A_{\text{stand}} \times W \times 2}$$

where:

$A_{\text{sample}}$  - Absorbance of sample was determined at 360 nm,

$A_{\text{stand}}$  - Absorbance of standard solution was determined at 360 nm,

C - Concentration of standard solution of rutin (g/ml),

W - weight (g) of sample for analyses,

2 - Volume (ml) of sample for analyses, 100 - Percent, %.

### Tannins assay

The analyses of tannins content in *Cotinus coggygia* were performed according to The International Pharmacopoeia and AOAC method, after modified methods. Measured 25 ml of this infusion into 1 L conical flask and add 25 ml indigo solution and 750 ml distilled deionized water (dd H<sub>2</sub>O). Titrated with 0.1 N water solution of KMnO<sub>4</sub> until blue solution changes to green, then add a few drops at time until solution becomes golden yellow. Was prepared standard solution of Indigo carmine (dissolve 6 g indigo carmine in 500 ml distilled deionized water (dd H<sub>2</sub>O) by heating, cool add 50 ml 96% - 98% H<sub>2</sub>SO<sub>4</sub>, diluted to 1 L and filter). For the blank similarly titrated mixture of 25 ml indigo carmine solution and 750 ml dd H<sub>2</sub>O. All samples were analyzed in duplicates [30].

Calculations

Calculations are based on averaging results from analyses of duplicate samples.

Calculate content (%) of tannins (T) in sample as follows:

$$T(\%) = \frac{(V - V_0) \times 0.004157 \times 250 \times 100}{g \times 26}$$

where:

V - Volume of 0.1 N water solution of KMnO<sub>4</sub> for titration of sample, ml;

V<sub>0</sub> - Volume of 0.1 N water solution of KMnO<sub>4</sub> for titration of blank sample, ml;

0.004157 - Tannins equivalent in 1 ml of 0.1 N water solution of KMnO<sub>4</sub>;

g – Mass of the sample for analyses, g;  
250 - Volume of volumetric flask, ml; 100 – Percent, %.

### 3 Statistical Analyses

All experiments were performed in triplicates. Analysis at every time point from each experiment was carried out in duplicate or triplicate. The statistical parameters are calculated in terms of the reproducibility of the experimental data using a statistical package universal ANOVA.

### 4 RESULTS AND DISCUSSION

Dry matter content in all experimental runs was determined and results were expressed on dry basis, which generally provides more accurate and reliable data comparison. The 50% ethanol extract of *Cotinus coggygia* showed the Table 1 and Table 2 of total phenols, total flavonoids, rutin and tannins in qualitative chemical analysis. Solid-liquid extraction was carried out by 50% aqueous ethanol for 120 min, which gave concentration of total polyphenols. The content for total phenolics and total flavonoids of *Cotinus coggygia* varied between 16.77 mg GAE/ml for 10 minutes and 24.11 mg GAE/ml for 120 minutes and 5.66 mg CE/ml for 10 minutes and 8.16 mg CE/ml for 120 minutes. This is shown in the (Table 1) using the gallic acid and catechin as standards. These results indicate that the higher antioxidant activity of the 50% ethanol extract of the *Cotinus coggygia* in 90 min and stopped in 120 min. than the 50% ethanol extract of *Cotinus coggygia* in 10 min may be in correlation with the phenolic and flavonoid contents of the extracts of the leaves.

TABLE 1

THE KINETIC VARIETIES OF 50% ETHANOL EXTRACT OF IN THE TOTAL PHENOLS AND TOTAL FLAVONOIDS

min	Total phenols, mg/ml dw	Total flavonoids, mg/ml dw
10	16.77±0.12 RSD 8.9% (n=3)	5.66±0.02 RSD 11.9% (n=3)
15	16.95±0.12 RSD 8.8% (n=3)	5.75±0.02 RSD 11.7% (n=3)
20	19.53±0.12 RSD 8.7% (n=3)	6.47±0.03 RSD 11.4% (n=3)
30	20.22±0.15 RSD 9.3% (n=3)	6.76±0.10 RSD 11.9% (n=3)
60	22.83±0.16 RSD 9.9% (n=3)	7.65±0.11 RSD 11.7% (n=3)
90	24.11±0.21 RSD 10.7% (n=3)	8.16±0.15 RSD 11.8% (n=3)
120	24.11± 0.21 RSD 10.7% (n=3)	8.16±0.16 RSD 11.9% (n=3)

The presence of rutin and tannins in 50% ethanol extract of *Cotinus coggygia* is significant. The content for rutin and tannins varied between 1.39 percent for 10 minutes and 2.26 percent for 120 minutes of rutin and tannins it was found that it was much higher than the content in 50% ethanol extract of *Cotinus coggygia* in 120 minutes between 10.45% and 7.33% in 10 minutes, respectively). The results were shown in the (Table 2), where the data was received using rutin as standard and Potassium permanganate as titrant. It is important to notice

that the comparison of the results for rutin and tannin contents in 50% ethanol extract of *Cotinus coggygia* will not be correct because of the different methods of analysis. The results and extraction curves indicated the exponential growth of extraction rate in the time for all examined particle classes.

TABLE 2

THE KINETIC VARIETIES OF 50% ETHANOL EXTRACT OF COTINUS COGGYGRIA IN RUTIN AND TANNINS

min	Rutin, %	Tannins, %
10	1.39±0.002 RSD 7.8; n=3	7.33±0.004 RSD 7.9; n=3
15	1.57±0.003 RSD 8.1; n=3	7.74±0.003 RSD 8.5; n=3
20	1.82±0.002 RSD 8.3; n=3	8.58±0.004 RSD 8.6; n=3
30	2.08±0.003 RSD 7.6; n=3	9.21±0.005 RSD 8.1; n=3
60	2.19±0.003 RSD 7.8; n=3	9.82±0.004 RSD 7.7; n=3
90	2.26±0.003 RSD 7.7; n=3	10.45±0.003 RSD 8.9; n=3
120	2.26±0.003 RSD 7.2; n=3	10.45±0.003 RSD 7.5; n=3

The kinetic study was performed by continuously measuring the absorbance of the extract using an UV-VIS spectrophotometer. The continuous measurement is faster and more accurate for kinetic studies of extraction compared to conventional discontinuous methods. In conventional methods, sampling is done manually at given intervals of time, which is not precise, as there is always a time gap between sampling and analysis, which contributes to errors during kinetic measurements.

In all experiments, the extraction yield was significantly time dependant and the profile clearly shows that the yield of *Cotinus coggygia* rises rapidly with time at first, and then less and less quickly as the progress of extraction continues. This behaviour can be attributed to the fact that during the initial stage of extraction, when the solvent penetrates into the solid, an extremely high concentration gradient is developed, resulting in high rates of mass transfer into the liquid phase. As the extraction time increases, the mass transfer of solutes from the solid phase to the fluid becomes more difficult, due to the decrease in concentration driving force between the solid and liquid phases. In addition, as the extraction time proceeds, the concentration of analytes in the solid phase decreases and both the solubility of the mixture and the extraction rate decrease simultaneously [31-33]. In all experiments, a higher extraction yield was observed especially from 10 to 90 min, with a lower yield seen from 90 to 120 min.

The WHO survey indicated that about 70–80% of the world's populations rely on non-conventional medicine, mainly of herbal source, for their primary healthcare [34, 35]. Plants are not only important to the millions of people to whom traditional medicine serves as the only opportunity for health care and to those who use plants for various purposes in their daily lives, but also as a source of new pharmaceuticals [36, 37]. So

far the extracts of many plant species have been examined for a number of biological activities, and their antimicrobial, anti-inflammatory, antioxidant, antimutagenic and cancer preventive effect have been partially described [37-41]. The use of a natural product with therapeutical properties has a long history. Plants are invaluable sources of pharmaceutical products [37, 42]. Many plant extracts have been used as a source of medicinal agents to cure urinary tract infections, cervicitis vaginitis, gastrointestinal disorders, respiratory diseases, cutaneous affections, helminthic infections and inflammatory process [37, 43, 44].

These medicinal plants are rich sources for naturally occurring antioxidants especially phenolic and flavonoids contents. These agents have ability to scavenge free radicals, super oxide and hydroxyl radicals, etc thus they enhance immunity and antioxidant defense of the body [45, 46]. Polyphenolic compounds like flavonoids, tannins, and phenolic acids, commonly found in such plants which contain various biological effects including antioxidant activity [47-49]. Mainly, phenolic compounds are very important for the free-radical scavenging and antioxidant activities of plants because these compounds react as hydrogen donors and thus neutralize the free-radicals [48-51]. Researchers have studied polyphenolic constituents having a potential to medicinal or nutraceutical properties including antioxidant activities [49, 52]. Therefore, the study of the importance and role of nonnutrient compounds, particularly phenolic acids, flavonoids and high molecular tannins as natural antioxidants has greatly increased [49, 53, 54]. There was a linear relationship between the antioxidant activities and phenolic compounds [49, 55, 56]. Different phytochemicals have various protective and therapeutic effects which are essential to prevent diseases and maintain a state of well being.

Phytochemical investigation of the ethanol extract of plant *C. coggygria* led to the isolation of several phenolic compounds [37, 57, 58]. Polyphenolic compounds are known to have antioxidant activity and it is likely that the activity of the extracts is due to these compounds. It is suggested that polyphenolic compounds have shown anticarcinogenic effects and potential to prevent cardiovascular and cerebrovascular diseases [37, 59].

Ethanol extract of *Cotinus coggygria* were analyzed for its phytoconstituents show that the dry leaves of *Cotinus coggygria* are rich in total phenols, total flavonoids, rutin and tannins to some extent (Table 1, Table 2). It is well that plant flavonoids and phenols in general, are the highly effective free radical scavenging and antioxidants. In this present study the phytochemical screening and quantitative estimation of the percentage of chemical constituents of the plants studied showed that the dry leaves of *Cotinus coggygria* were rich in rutin and tannins.

*Cotinus coggygria* was the most active species among the plants selected for the screening. This plant is largely used in the Balkan and Anatolian regions to cure wounds and reduce inflammations, as well as for the treatment of gastrointestinal and respiratory disorders [60, 61]. In Asiatic countries, *C. coggygria* is also known as a bactericide and frequently administered against hepatitis and even anemia [61, 62]. A relative of

this species, *Rhus coriaria*, which grows in the Mediterranean region, has demonstrated inhibitory properties towards *Streptococcus mutans* and *S. sanguinis*, common components of dental plaque [61, 63]. These authors attributed this effect to the presence of large amounts of tannins in the plant. Tannins can then generate smaller phenolics compounds (pyrogallol, catechol, and ellagic acid) with known bactericidal actions. Similarly, *C. coggygria* is very rich in phenolic compounds [61, 64] and displays a significant antimicrobial activity.

Tannins are polyphenolic substances with various molecular weights and a variable complexity [30, 49, 65]. Tannins exhibit many biologically significant functions, such as protection against oxidative stress, and degenerative diseases [30,49]. Rutin is the glycoside between the flavonol quercetin and the disaccharide rutinose [29, 49]. Rutin is one of the bioactive flavonoid compounds, which are present in substantial amounts in plants.

Phenolic compounds have multiple biological effects, such as anti-atherogenic, antioxidant, anti-inflammatory, cardioprotective, antimicrobial, anticarcinogenic and neuroprotective [66-68]. The contents of total phenolics, tannins and flavonoids of the *C. coggygria* extract and its derived fractions are given in Table 1 and 2. Plant secondary metabolites such as terpenoids, phenolic compounds, alkaloids and lectins exert an antimicrobial effect [68, 69].

Many efforts have been made to discover new antimicrobial compounds from various sources such as animals, microorganisms and plants. Plants possess antimicrobial natural products to protect themselves [37, 69, 70]. Antimicrobial activities of various herbs and spices in plant leaves, flowers, stems, roots or fruits have been reported [37,71- 74].

Microbial antibiotic resistance is increasing worldwide and can be demonstrated against all classes of antibiotics. Moreover, multi-drug-resistant strains have emerged in many species of pathogenic microorganisms [68, 75]. Consequently, there is a growing interest in new sources of antimicrobial agents. The phenolic compounds are not only responsible for the antimicrobial activity of *C. coggygria*. This is in agreement with the previous demonstration of the high antimicrobial effect of the essential oil from *C. coggygria* leaves on Gram-positive bacteria and fungi [8, 68].

The antioxidant activity of phenolic compounds is based on their ability to donate hydrogen atoms or electrons, chelate metal cations and scavenge free radicals [68, 76, 77]. The process of LP has an important role in oxidative stress development. The peroxidation of unsaturated lipids of the cell membrane leads to cell membrane damage, loss of membrane integrity and consequently to cell and tissue damage [68, 78].

ROS damage cell membranes and DNA, causing potentially cancerous mutations. They are also implicated in inflammatory processes [68, 79]. Phenolic compounds have an ability to inhibit the production of ROS or to neutralise radicals that are involved in the inflammatory process as signalling molecules [68, 80].



## 5. CONCLUSION

In conclusion, the results of this research showed that total phenolic, total flavonoid, rutin and tannin contents are important components in 50% ethanol extract of Bulgarian dry leaves of *Cotinus coggygia* this plant, and some of the pharmacological effects. Results of the present work clearly shows that the yield of *Cotinus coggygia* rises rapidly with time at first, and then less and less quickly as the progress of extraction continues. The Bulgarian dry leaves of *Cotinus coggygia* are an important component, providing some protective/preventative health effect.

## Acknowledgement

We are highly thankful to Physics Department, Faculty of Science, University of Tabuk, Saudi Arabia and National Centre of Public Health and Analyses, Department "Public Health and Health risk", Akad, Sofia, Bulgaria for keen support and help in our present research work [81-83].

## REFERENCES

- [1] Singh P, Shrivastava R, Saxena RC, Sharma M, Karchuli MS, Tripathi J. Phytochemical screening and evaluation of antioxidant activity of *Parkinsonia aculeata* L. Leaves extract. Int J Pharm Tech Res 2011;3:1952-7
- [2] Abha Shukla, Swati Vats, Rishi Kumar Shukla, Proximate Composition, Nutritive Value and Evaluation of Antioxidant Potential of Stem of *Dracaena Reflexa* LAM, Int J Pharm Pharm Sci, 2014 Vol 6, Issue 11, 360-364
- [3] Janifer X Raj, Chaurasia AP, Vajpayee PK, Pal Murugun M, Singh Shashibala. Antioxidant activity and phytochemical investigation on a high altitude medicinal plant *Dracocephalum heterophyllum* benth, Pharmacogn J 2010;2:112-7
- [4] Gurav Shailendra, Deshkar N, Gulkari V, Durgakar N, Patil A. Free radical scavenging activity of *Polygala chinensis*. Pharmacol 2007;2:245-53
- [5] Jayaprakasha, G. K., Selvi, T., & Sakariah, K. K. (2003). Antibacterial and antioxidant activities of grape seed extracts. Food Research International, 36, 117-122.
- [6] Ana Bucić-Kojić, Mirela Planinić, Srećko Tomas, Mate Bilić, Darko Velić, Study of solid-liquid extraction kinetics of total polyphenols from grape seeds, Journal of Food Engineering 81 (2007) 236-242
- [7] Atanassova M, Georgieva S, Ivancheva K. Total phenolic and total flavonoids contents, antioxidant capacity and biological contaminants in medicinal herbs. J Univ Chem Techn Metal 2011;46:81-8
- [8] Novakovic M, Vučkovic I, Janackovic P, Sokovic M, Tesevic V and Milosavljevic S (2007) composition, antibacterial and antifungal activity of the essential oils of *Cotinus coggygia* from Serbia. J Serb Chem Soc 72:1045-1051
- [9] Sanja Matic, Snezana Stanic, Desanka Bogojevic, Slavica Solujic, Nevena Grdovic, Melita Vidakovic and Mirjana Mihailovic, (2011) Genotoxic potential of *Cotinus coggygia* Scop. (Anacardiaceae) stem extract in vivo. Genetics and Molecular Biology, 34, 2, 298-303
- [10] Landzhev Chemical, I., 2010. Encyclopedia of Medicinal Plants in Bulgaria. Trud, Sofia, 551 pp. (Bg).
- [11] D. Ivanova, D. Pavlov, M. Eftimov, K. Kalchev, M. Nashar, M. Tzaneva and S. Valchev Subchronic Toxicity Study of Ethanol Infusion from *Cotinus Coggygia* Wood in Rats, Bulgarian Journal of Agricultural Science, 19 (2) 2013, 182-185 Agricultural Academy
- [12] Rayne S and Mazza G (2007) Biological activities of extracts from Sumac (*Rhus spp.*). Plant Foods Hum Nutr 62:165-175.
- [13] Demirci B, Demirci F and Baser KH (2003) Composition of the essential oil of *Cotinus coggygia* (Scop.) from Turkey. Flavour Fragr J 18:43-44.
- [14] Ivanova D, Geroval D, Chervenkov T and Yankova T (2005) Polyphenols and antioxidant capacity of Bulgarian medicinal plants. J Ethnopharmacol 96:145-150.
- [15] Heim, K. E., Tagliaferro, A. R., & Bobilya, D. J. (2002). Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. The Journal of Nutritional Biochemistry, 13, 572-584
- [16] Yilmaz, Y., & Toledo, R. T. (2004). Major flavonoids in grape seeds and skins: antioxidant capacity of catechin, epicatechin, and gallic Acid. Journal of Agricultural and Food Chemistry, 52, 255-260.
- [17] Baydar, N. G., Özkan, G., & Sağdıç, O. (2004). Total phenolic contents and antibacterial activities of grape (*Vitis vinifera* L.) extracts. Food Control, 15, 335-339.
- [18] Lapornik, B., Prošek, M., & Golc Wondra, A. (2005). Comparison of extracts prepared from plant by-products using different solvent and extraction time. Journal of Food Engineering, 71, 214-222.
- [19] Pekić, B., Kovač, V., Alonso, E., & Revilla, E. (1998). Study of the extraction of proanthocyanidins from grape seeds. Food Chemistry, 61(1/2), 201-206
- [20] Palma, M., & Taylor, L. T. (1999). Extraction of polyphenolic compounds from grape seeds with near critical carbon dioxide. Journal of Chromatography A, 849, 117-124.
- [21] Bandonienė, D., Pukalskas, A., Venskutonis, P. R., & Gruzdienė, D. (2000). Preliminary screening of antioxidant activity of some plant extracts in rapeseed oil. Food Research International, 33, 785-791.
- [22] Moure, A., Cruz, J. M., Franco, D., Domínguez, J. M., Sineiro, J., Domínguez, H., et al. (2001). Natural antioxidants from residual sources. Review. Food Chemistry, 72, 145-171.
- [23] Škerget, M., Kotnik, P., Hadolin, M., Rižner Hraš, A., Simonić, M., & Knez, Ž. (2005). Phenols, proanthocyanidins, flavones and flavonols in some plant materials and their antioxidant activities. Food Chemistry, 89, 191-198.
- [24] Guendez, R., Kallithraka, S., Makris, D. P., & Kefalas, P. (2005). Determination of low molecular weight polyphenolic constituents in grape (*Vitis vinifera* sp.) seed extract: correlation with antiradical activity. Food Chemistry, 89, 1-9.
- [25] Tsao, R., & Deng, Z. (2004). Separation procedures for naturally occurring antioxidant phytochemicals. Review. Journal of Chromatography B, 812, 85-99.
- [26] Pinelo, M., Del Fabbro, P., Manzocco, L., Nuñez, M. J., & Nicolini, M. C. (2005). Optimization of continuous phenol extraction from *Vitis vinifera* by products. Food Chemistry, 92, 109-117.
- [27] Shi, J., Yu, J., Pohorly, J., Young, J. C., Bryan, M., & Wu, Y. (2003b). Optimization of extraction of polyphenols from grape seed meal by aqueous ethanol solution. Food Agriculture and Environment, 1(2), 42-47.
- [28] Marinova D., Ribarova F., Atanassova M. Total phenolics and total flavonoids in Bulgarian fruits and vegetables, Journal of the University of Chemical Technology and Metallurgy. 2005. Vol. 40, N 3. P. 255-260.

- [29] Atanassova M and Bagdassarian V. 2009. Rutin content in plant products. *Journal of the University of Chemical Technology and Metallurgy* 44(2): 201-203.
- [30] Atanassova M., Christova-Bagdassarian V. (2009). Determination of tannins content by titrimetric method for comparison of different plant species. *Journal of the University of Chemical Technology and Metallurgy*, 44, 4, pp. 413-415.
- [31] Goula, A. M. (2013). Ultrasound-assisted extraction of pomegranate seed oil – Kinetic modeling. *Journal of Food Engineering*, 117, 492–498.
- [32] Hatami, T., Cavalcanti, R. N., Takeuchi, T. M., & Meireles, M. A. A. (2012). Supercritical fluid extraction of bioactive compounds from Macela (*Achyrocline satureioides*) flowers: Kinetic, experiments and modeling. *The Journal of Supercritical Fluids*, 65, 71–77.
- [33] Mahesha M. Poojary, Paolo Passamonti, Extraction of lycopene from tomato processing waste: Kinetics and modelling, *Food Chemistry* 173 (2015) 943–950
- [34] Chan K, Some aspects of toxic contaminants in herbal medicines. *Journal of Chemosphere*. 2003; 52:1361–1371.
- [35] Rajeev Singh, Pawan Kumar Verma, Gagandeep Singh, Total phenolic, flavonoids and tannin contents in different extracts of *Artemisia absinthium*, *Journal of Intercultural Ethnopharmacology* 2012; 1(2):101-104
- [36] De Boer H J, Kool A, Broberg A, Mziray W R, Hedberg I, Levenfort J J 2005 Anti-fungal and anti-bacterial activity of some herbal remedies from Tanzania. *J Ethnopharmacol* 96 (3): 461–469
- [37] Sanja Matić, Snežana Stanić, Slavica Solujić, Tanja Milošević, Neda Niciforović, Biological properties of the *Cotinus coggygria* methanol extract, *Periodicum Biologorum*, Vol. 113, No 1, 87–92, 2011
- [38] Baricevis D, Bartol T 2000 The biological/pharmacological activity of the *Salvia* genus. In: Kintzois S E (ed) *Pharmacology*. Sage. The Genus *Salvia*. Haewood Academic Publishers, Amsterdam, p 143–184
- [39] Mitić D, Vukicević-Gačić B, Knezević-Vukcević J, Berić T, Nikolić B, STanković S, Simić D 2001 Natural antioxidants and their mechanisms in inhibition of mutagenesis. In: Kreft M, Skrbanja V (eds) *Molecular and Genetic Interactions Involving Phytochemicals*. Univ. Ljubljana and Slovenian, Academy of Sciences and Arts. Ljubljana, p 67–74
- [40] Vujošević M, Blagojević J 2004 Antimutagenic effect of extract from sage (*Salvia officinalis*) in mammalian system in vivo. *Acta Vet Hung* 52: 439–443
- [41] Faried A, Kurnia D, Faried L S, Usman N, Miyazaki T, Katov H, Kuwano H, 2007 Anticancer effects of gallic acid isolated from Indonesian herbal medicine, *Phaleria macrocarpa* (Scheff.) Boerl on human cancer cell lines. *Int J Oncol* 30: 605–613
- [42] Olalder J A 2005 The systemic theory of living systems and relevance to CAM. Part I: the theory. *Evid Based Complement Alternat Med* 2: 13–18
- [43] Bratner A, Grein E 1994 Antibacterial activity of plant extracts used externally in traditional medicine. *J Ethnopharmacol* 44: 35–40
- [44] Meyer J J M, Afolayan A J, Taylor M B, Engelbrecht L 1996 Inhibition of herpes simplex virus type 1 by aqueous extract from shoots of *Helichrysum aureonitens* (Asteraceae). *J Ethnopharmacol* 52: 41–43
- [45] Atoui K, Mansouri A, Bosku G, Kefalas P. Tea and herbal infusions: their antioxidant activity and phenolic profile. *Food Chem*. 2005; 89: 27-36. =12
- [46] Kahkonen M P, Hopia A I, Vuorela H J, Rauha J P, Pihlaja K, Kujala T S and Heinonen M. 1999. Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agriculture and Food Chemistry* 47: 3954-3962.=13
- [47] Mohammad N, Majumder M S, Rashed-al-Qayum, Khan M D, Bhattacharjee S and Kar A. 2013. Phytochemical screening of medicinal plant-mikania cordifolia and determination of its characteristics. *Mintage Journal of Pharmacy and Medical Sciences* 2(1): 14-17.=14
- [48] Tanaka T, Kuie C W, Nagashima Y and Taguchi. 1988. Applications of antioxidative maillard reaction products from histidine and glucose to sardine products. *Nippon Suisan Gakkaishi* 54: 1409-1414.=15
- [49] Valentina L. Christova-Bagdassarian, Kristine S Bagdassarian, Maria S Atanassova and M Ayaz Ahmad, Comparative Analysis of Total Phenolic and Total Flavonoid Contents, Rutin, Tannins and Antioxidant Capacity in *Apiaceae* and *Lamiaceae* families, *Indian Horticulture Journal*; 4(3/4): 131-140, July-December (2014)=16
- [50] Kulisic T, Radonic A, Katalinic V and Milos. 2004. Use of different methods for testing antioxidative activity of oregano essential oil. *Food Chemistry* 85(4): 633-640.=17
- [51] Siddhuraju P. 2006. The antioxidant activity and free radical scavenging capacity of phenolics of raw and dry heated moth bean (*Vigna ac-onitifolia*) (Jacq.) Marechal seed extracts. *Food Chemistry* 99: 149-157.=18
- [52] Siddhuraju P and Becker K. 2007. The antioxidant and free radical scavenging activities of processed cowpea (*Vigna unguiculata* (L.) Walp.) seed extracts. *Food Chemistry* 101: 10-19.=19
- [53] Sahoo H B, Bhattaisra S K, Biswas U K and Sagar R. 2013. Estimation of total phenolics and flavonoidal contents as well as *in vitro* antioxidant potential of *Apium leptophyllum* Pers. *Herba Polonica* 59(3): 37-50.=20
- [54] Gao X, Ohlander M, Jeppsson N, Bjork L and Trajkovski V. 2000. Changes in antioxidant effects and their relationship to phytonutrients in fruits of sea buckthorn (*Hippophae rhamnoides* L.) during maturation. *Journal of Agriculture and Food Chemistry* 48: 1485-1490.=21
- [55] Huang B, Ban X, He J S, Zeng H, Zhang P and Wang Y W. 2010. Hepatoprotective and antioxidant effects of the methanolic extract from *Halenia elliptica*. *Journal of Ethnopharmacology* 131: 276-281.=22
- [56] Atanassova M and Georgieva S. 2010. Comparative polyphenol composition and antioxidant capacity of the Bulgarian plants (Dry Herbs). *EJEAF Che* 9(9): 1514-1523.=23
- [57] Stathopoulou K, Magitis P, Karapanigiotis I, Valianou L, Chrysosoulakis Y 2007 Phytochemical analysis of *Cotinus coggygria* heartwood. Identification of isolated colorants in historical art objects. In: Proceedings of the 55th International Congress and Annual Meeting of the Society for Medicinal Plant Research, Austria. *Planta Med* 73: 163
- [58] Zdunić G, Stević T, Šavikin K, Menković N, Janković T 2007 Antimicrobial activity of some medicinal plants growing in Serbia and Montenegro. In: Proceedings of the 55th International Congress and Annual Meeting of the Society for Medicinal Plant Research, Austria. *Planta Med* 73: 150
- [59] KOondratyuk P, Pezzuto J M 2004 Natural product polyphenols of relevance to human health. *Pharm Biol* 42: 46–63
- [60] K. L. Ploetz, *An Ethnobotanical Study of Wild Herb Use in Bulgaria*, Michigan Technological University, 2000.
- [61] Gianmaria F. Ferrazzano, Lia Roberto, Maria Rosaria Catania, Angela Chiaviello, Antonino De Natale, Emanuela Roscetto, Gabriele Pinto, Antonino Pollio, Aniello Ingenito, and Giuseppe Palumbo, Screening and Scoring of Antimicrobial and Biological Activities of Italian Vulnerary Plants against Major Oral Pathogenic Bacteria, *Hindawi Pub-*

- lishing Corporation Evidence-Based Complementary and Alternative Medicine Volume 2013, Article ID 316280, 10 pages
- [62] M. Ajaib, Z.-U. Khan, N. Khan, and M. Wahab, "Ethnobotanical studies on useful shrubs of District Kotli, Azad Jammu & Kashmir, Pakistan," *Pakistan Journal of Botany*, vol. 42, no. 3, pp. 1407–1415, 2010.
- [63] E. Babpour, S. A. Angaji, and S. M. Angaji, "Antimicrobial effects of four medicinal plants on dental plaque," *Journal of Medicinal Plant Research*, vol. 3, no. 3, pp. 132–137, 2009.
- [64] D. Ivanova, D. Gerova, T. Chervenkov, and T. Yankova, "Polyphenols and antioxidant capacity of Bulgarian medicinal plants," *Journal of Ethnopharmacology*, vol. 96, no. 1-2, pp. 145–150, 2005.
- [65] Makkar H P S. 2003. Effects and fate of tannins in ruminant animals, adaptation to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Ruminant Research* 49: 241-256.=24
- [66] Manach C, Mazur A, Scalbert A. 2005. Polyphenols and prevention of cardiovascular diseases. *Curr Opin Lipidol* 16(1): 77–84.
- [67] Middleton E, Kandaswami C, Theoharides TC. 2000. The effect of plant flavonoids on mammalian cells: implication for inflammation, heart disease and cancer. *Pharmacol Rev* 52(4): 673–751.
- [68] Mirjana Marčetić, Dragana Božić, Marina Milenković, Nevenka Malešević, Siniša Radulović and Nada Kovačević, Antimicrobial, antioxidant and anti-inflammatory activity of young shoots of the smoke tree, *Cotinus coggygria* Scop, *Phytother. Res.* 27: 1658–1663 (2013)
- [69] Cowan MM. 1999. Plant Products as Antimicrobial Agents. *Clin Microbiol Rev* 12: 564–582.
- [70] Tomoko N, Tabachi A, Hiromu T, Yuka I, Hiroko M, Muneakaju I, Totshiyki T, Tetsuro I, Fujio A, Irija I, Tsutomu N, Kazuhito W 2002 Antibacterial activity of extracts prepared from tropical and subtropical plants on methicillin-resistant *Staphylococcus aureus*. *J Health Sci* 48: 273–276
- [71] Rajeh M A B, Zuraini Z, Sasidharan S, Latha Y L, Amutha S 2010 Assessment of *Euphorbia hirta* L. leaf, flower, stem and root extracts for their antibacterial and antifungal activity and brine shrimp lethality. *Molecules* 15(9): 6008–6018
- [72] Wamidh H T, Adel M M 2010 Antimicrobial, cytotoxicity and phytochemical screening of Jordanian plants used in traditional medicine. *Molecules* 15(3): 1811–1824
- [73] Sumathi P, Parvathi A 2010 Antimicrobial activity of some traditional medicinal plants. *J Med Plants Res* 4(4): 316–321
- [74] Afolabi M O, Adegoke G O, Mathooko F M, 2011 Phytochemical characterization of the extracts of *Aframomum daniellii* flower, leaf, stem and root. *Afr J Agr Res* 6(1): 97–101
- [75] Orhan DD, Özçelik B, Özgen S, Ergun F. 2010. Antibacterial, antifungal, and antiviral activities of some flavonoids. *Microbiol Res* 165(6): 496–504.
- [76] Bondet V, Brand-Williams W, Berset C. 1997. Kinetics and Mechanisms of Antioxidant Activity using the DPPH<sup>•</sup> Free Radical Method. *Lebensm-Wiss u-Technol* 30: 609–615.
- [77] Afanas'ev IB, Dorozhko AI, Brodskii AV, Kostyuk VA, Potapovitch AI. 1989. Chelating and free radical scavenging mechanisms of inhibitory action of rutin and quercetin in lipid peroxidation. *Biochem Pharmacol* 38: 1763–1769.
- [78] Halliwell B. 2006. Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. *Plant Physiol* 141 (2): 312–322.
- [79] Roessner A, Kuester D, Malfertheiner P, Schneider-Stock R. 2008. Oxidative stress in ulcerative colitis-associated carcinogenesis. *Pathol Res Pract* 204(7): 511–524.
- [80] Santangelo C, Vari R, Scazzocchio B, Di Benedetto R, Filesi C, Masella R. 2007. Polyphenols, intracellular signalling and inflammation. *Ann Ist Super Sanita* 43: 394–405.
- [81] Anghel Drugarin Cornelia Victoria, M. Ayaz Ahmad, N. Ameer Ahmad, Draghic Silviu. (2015). The Mathematical Study of Data Transmission in Digital Electronics. *International Journal of Advanced Research (IJAR)*. 3(3): 697- 702.
- [82] Anghel Drugarin Cornelia Victoria, M. Ayaz Ahmad, N. Ameer Ahmad, Vyacheslav V. Lyashenko. (2015). Algorithmic Research and Application Using the Rayleigh Method", *International Journal of Science & Research (IJSR)*. 4(4): 1669-1671.
- [83] Dragos Pasculescu, Remus Dobra and M. Ayaz Ahmad (2016). Dosimetric Quantity System for Electromagnetic Fields Bio-effects" , *International Journal of Scientific Research*, Vol. 5(2), (2016), 26-32.