

X-ray Satellites Spectra in the $L\alpha_1$ Region of 4d Transition Elements

Dr. Sameer Sinha, Vinay Kumar Pandey, Ajay Vikram Singh

Abstract- We have used Plasmon theory to explain Energy Satellites and relative intensity of high energy X-ray satellites α_3 , α_4 & α_5 with respect to $L\alpha_1$ parent line in 4d Transition Metal (Zr, Nb, Mo, Ru, Rh) and estimated values are in agreement with the calculated values of Surendra Poomia and S.N.Soni.

Keywords - $L\alpha_1$ X-ray satellites, M vacancy $L\alpha_1$ transitions, X-Ray emission spectra.

INTRODUCTION

In the characteristic X-ray Spectra, Diagram as well as non Diagram lines are present. Those lines which fit in the conventional energy level diagram are called Diagram lines. & those lines which do not fit in the conventional energy level diagram are called non diagram lines. It is also known as "Satellites or Second order lines". Satellites are generally of weak intensity lines & are found close to more intense parent line. The satellites which are observed on higher energy side are called high energy satellites (HES) whereas those are observed on lower energy side are called lower energy satellites (LES). First Siegbahn & Stenstroem observed these satellites in the K-Spectra of element from Cr (24) to Ge (32) while Coster Theraeus & Richtmyer in the L-Spectra of element from Cu (29) to Sb (51) & Hajlmar, Hindberg & Hirsch in the M-Spectra of elements from Yb (70) to U (92). Several theories were proposed from time to time to explain the origin of these satellites. Out of these theories the plasmon theory is found to be the most suitable theory especially for those satellites.

Plasmon theory was first proposed by Bohm & Pines which are extended by Houston, Ferrel, Noziers & Pines. According to this theory the low energy plasmon satellites are emitted when valence electron excites a plasmon during the annihilation of core hole conversely if Plasmon pre exists, its energy add up to the energy of diagram line.

* Reader, Ganpat Sahai Post Graduate College, Sultanpur, U.P. India

** Lecturer, Ganpat Sahai Post Graduate College, Sultanpur, U.P. India

*** Associate Professor, Rajarshi Rananjay Singh Institute of Management & Technology, Amethi, CSJ Nagar, U.P., India
E-Mail ID –

*** ajay_gspgcs@rediffmail.com

** vinay_gspg@rediffmail.com

The radiation less reorganization of electronic shell of an atom is known as Auger effect. Auger satellites have also been observed by Korbar and Mehlhorn [1] Haynes et al. [2] Edward and Rudd [3]. Theoretical explanation for K series Auger spectrum was given by Burhop and Asaad [4] using intermediate coupling. Later on more refined theory, using relativistic and configuration interaction has been used by Listengarter [5] and Asaad [6]

In Auger primary spectra, one can also observe secondary electron peaks close to the primary peaks are produced by incident electrons which have undergone well energy losses. The most common source of such energy loss in the excitation of collective plasma oscillations of the electrons in the solid. This gives rise to a series of plasma peaks of decreasing magnitude spaced by energy $\hbar\omega_p$ where ω_p is the frequency of plasma oscillation.

Auger peaks are also broadened by small energy losses suffered by the escaping electrons. This gives rise to a satellite on the low energy of the Auger peak. Energy loss peaks have well defined energy with to primary energy.

The involvement of Plasmon oscillation in the X-ray emission or absorption spectra of solids has been widely studied during the last few decades and has been recognized that the electron-electron interaction has played an important role.

This Paper is devoted to Plasmon theory to explain the Energy Satellites and relative intensity of high energy X-ray satellites with respect to $L\alpha_1$ parent line in 4d Transition Metal (Zr, Nb, Mo, Ru, Rh, Rd) and estimated values are in agreement with the calculated values of Surendra Poomia and S.N.Soni.

According to Plasmon theory, if the valence electron, before filling the core vacancy, also excites a Plasmon, then the energy $\hbar\omega_p$ needed for the excitation of Plasmon oscillation is taken from the transitioning valence electron so that the emitted radiation will be derived off an energy $\hbar\omega_p$ and a low energy satellites will emitted whose separation from the main X-ray line will correspond to $\hbar\omega_p$. On the other hand if the Plasmon pre exists, during the X-ray

emission process, then, on its decay it can give its energy to the transiting valence electron before it annihilates the core vacancy. Thus the energy of emitted X-ray photon will be higher than the main emission line and by an amount $\hbar\omega_p$ giving rise to high energy satellite.

Surendra Poonia and S.N.Soni have observed low and high energy satellite peaks in 4d Transition Metal (Zr, Nb, Mo, Ru, Rh, Rd). A close approximation of their tables shows that some satellites are at a distance of $\hbar\omega_p$ (Plasmon energy) from the main emission line. This observation forced us to think that these might be due to Plasmons emission and absorption.

MATHEMATICAL CALCULATION –

In order to confirm the involvement of Plasmon in the emission of X-ray satellites the relative intensity of single Plasmon satellites must be calculated. In this process first we deal with mathematical details of canonical transformation carried out over the model Hamiltonian of the system. Thus the energy separation ΔE of the low and high energy Plasmon satellite from the corresponding main line should be equal to the quantum of Plasmon energy $\hbar\omega_p$ which is given by [10]

$$\Delta E = \hbar\omega_p = 23.8 \sqrt{\left(\frac{Z \cdot \sigma}{w}\right)} \text{ev} \quad 1$$

Where Z = No. of unpaired electrons taking part in plasma oscillation
 σ = Specific gravity
 w = Molecular Weight

This equation can be derived as given below. From the classical consideration, we get the frequency of Plasmon oscillation as

$$\omega_p = \left(\frac{4\pi n e^2}{m}\right)^{1/2} \quad 2$$

Hence the amount of energy given to Plasmon becomes

$$E_p = \hbar\omega_p = h \left(\frac{4\pi n e^2}{m}\right)^{1/2}$$

In this equation we can write

$$n = \frac{L\sigma Z}{w}, \quad Z \text{ and } W \text{ are defined}$$

above and L is the Avogadro number. By putting the numerical value of constant, we get the Plasmon energy as

$$\Delta E = \hbar\omega_p = 23.8 \sqrt{\left(\frac{Z \cdot \sigma}{w}\right)} \text{ev} \quad 3$$

Our calculated values of ΔE have been compared with the Scrocco's experimental value. And We have also calculated the relative intensity of plasmon satellites, which is different in different processes. If the excitation of plasmon occurs during the transport of the electron through the solid, it is known as

extrinsic process of plasmon excitation. The plasmon can also be excited by another method known as intrinsic process. In this process, excitation of plasmon takes place simultaneously with creation of a hole. Bradshaw et al have further divided core hole excitation into two classes,

1 - Where the number of slow electrons are conserved.

2 - Where the number of slow electrons are not conserved

The Author has calculated relative intensity in both the cases with new modification in the light of Bradshaw [11] and Lengreth [12] work, which explains that not only intrinsic process but extrinsic process and their relative contribution may also contribute in relative intensities. The combined effect of intrinsic and extrinsic plasmon excitation intensity variation was also suggested by various Authors [12, 15, 16, 17 18] as:

$$i = \frac{I_s}{I_m} = \alpha^n \sum_{m=0}^n \frac{\left(\frac{\beta}{\alpha}\right)^m}{m!} \quad 4$$

The value of β is taken as $\beta = 0.12r_s$, which is purely intrinsic, $r_s = (47.11 / \hbar w_s)^{2/3}$ is dimensionless parameter and $\alpha = 0.47 r_s^{1/2}$ in the place of $\alpha = (1+L/L)^{-1}$ used by Pardee et. al.(13). The equation (3) contains a series of terms. The first term of the equation is purely extrinsic, while second term is purely intrinsic. The other terms are containing the relative contributions of both extrinsic and intrinsic. The specialty of this formula is that each term alone or simultaneously with other terms is able to give the relative intensity. This formula also includes both the categories mentioned by Bradshaw and gives better results as compared than traditional methods for calculation of the relative intensity. Using the values of α , β and r_s in equation (4)

Using the equation (4), the author has for the first time, calculated the relative intensity of high energy X-ray satellites with respect to $L\alpha_1$ parent line in 4d Transition Metal (Zr, Nb, Mo, Ru, Rh, Rd) metals. Our calculated and estimated values are in agreement with the calculated values of Surendra Poonia and S.N.Soni [14].

Reference

1. Korbar H. & Mehlhorn W.A. ; Phys. 191, (1966) 217.
2. Haynes S.K. & Velinsky, M & Velinsky L.J. ; Nucl. Phys. A99 (1967), 537.
3. Rudd M.E. & Edward & Volz, D.J. ; Phys Rev. 151, (1966), 28.
4. Asaad, W.N. & Burhop E.H.S. ; Proc. Phys. Soc., London 71, (1958), 369.
5. Listengarten, M.A. ; Bull Acad. Sci. U.S.S.R., Phys. Ser. 26 (1962), 182.
6. Asaad, W.N. ; Nucl. Phys. 66, (1965b), 494.

7. M.Scrocco in photoemission spectra of Pb.(II) halide; Phys. Rev. B25 (1982) 1535-1540 .
8. M.Scrocco , Satellites in X-ray Photo electron spectroscopy of insulator I 32 (1985) 1301-1306
9. M.Scrocco , Satellite in X-ray Photo electron spectroscopy of insulators II 32 (1985) 1307-1310
10. L.Marton , L.B.Lader and H. Mendlowitz; Adv. Electronic and Electro Physics; edited by L.M arton Academic , New York 7 (1955) , 225
11. A. M. Bradshaw, Cederbaurn S.L,Domeke W. & Krause Jour. Phys C: Solid State **Phys. 7, 4503, 1974**
12. D. C. Lengreth, Phys. Rev. Letter, 26, 1229, 1971
13. W. J. Pardee, G.D. Mahan, D. E. Eastman, R.A. Pollak, L. Ley, F.R. McFeely, S.P. Kowalczyk and D.A. Shirely, Phys. Rev. B, 11, 3614, 1975.
14. Surendra poonia and S.N.Soni , Indian journal of pure and applied physics , vol.45, feb.2007 pp-119-126
15. S. K. Srivastava, Amar Bahadur, *Fizika A*, 17, 117, 2008
16. S. K. Srivastava Leonardo Journal of Sciences , 13 , 39 , 2008.
17. Sameer Sinha & Ajay Vikram Singh , International Journal of Scientific & Engineering Research, Volume 2, Issue 2, April -2011
18. Sameer Sinha & Ajay Vikram Singh , International Journal of Scientific & Engineering Research, Volume 2, Issue 2, February-2011

S.No.	NAME	SYMBOL	Z	W	Sp.Gravity	Exper. Value [14]	Author Value
1	ZIRCONIUM	Zr(40)	1	91.224	6.51	6.54	5.44
2	NIBOLIUM	Nb(41)	1	92.906	8.58	6.51	6.19
3	MOLYBDENUM	Mo(42)	1	95.94	10.28	6.54	6.67
4	RUTHENIUM	Ru(44)	1	101.07	12.45	6.85	7.15
5	RHODIUM	Rh(45)	1	102.91	12.41	7.36	7.07
6	PALLADIUM	Pd (46)	1	106.42	12.02	7.59	6.84

Energy separation ΔE at $L\alpha_1$ satellite of 4-d Transition element

Relative Intensity of $L\alpha_1$ satellite of 4-d Transition element

S.No.	Name of Metal	Surface Energy	Rs	Alpha	Beta	Author intensity	Exp. Intensity [14]	Symbol
1	ZIRCONIUM	5.44	4.22	0.97	0.51	0.78065	0.780012	$\beta+0.1+\beta^2/2\alpha$
2	NIBOLIUM	6.19	3.87	0.92	0.46	0.7780	0.770182	$\beta+0.1+\beta^2/2\alpha$
3	MOLYBDENUM	6.67	3.68	0.90	0.44	1.3530	1.368272	$2*(\beta+0.1+\beta^2/2\alpha)$
4	RUTHENIUM	7.15	3.51	0.88	0.42	0.7834	0.7434	$2*\beta - 0.1$
5	RHODIUM	7.07	3.54	0.88	0.42	0.8655	0.8498	$2*\beta$
6	PALLADIUM	6.84	3.77	0.91	0.45	0.8623	0.9044	$2*\beta$

